

INFLUENCE OF PREY TYPES ON CERTAIN BIOLOGICAL CHARACTERISTICS OF *Hippodamia tridecimpunctata* L. UNDER CONSTANT TEMPERATURE AND RELATIVE HUMIDITY.

Mohamed, Nadia E.¹ ; A. A. Ghanim² and A. A. Saleh³

^{1,3} Plant Protection Res. Inst., Agric. Res. Center, Dokki Giza, Egypt.

² Economic Entomology Dept., Fac. Agric., Mansoura University, Egypt.

ABSTRACT

Laboratory experiments were carried out to study certain biological characteristics of the coccinellid predator, *Hippodamia tridecimpunctata* L. when reared on the following aphid species; *Sitobion avenae* Fabr.; *Rhopalosiphum maidis* (Fitch); *Acyrtosiphon pisum* (Harris) and *Aphis gossypii* Glover under constant temperature and relative humidity.

The data revealed that the total developmental time from egg hatching to adult eclosion of the coccinellid predator was differed significantly when reared on the four tested aphid species. Mortality percentage of *H. tridecimpunctata* from egg hatching to adult emergence ranged from 4.98 % when fed on *A. pisum* to 9.05% on *S. avenae*.

Based on statistical analysis, the mortality percentage showed significant difference among the four aphid species. The total consumption rate per *H. tridecimpunctata* larva from the four aphid species were 695.80 ± 10.75 ; 781.83 ± 15.68 ; 612.50 ± 9.55 and 894.53 ± 11.61 aphid individuals when reared on *S. avenae* ; *R. maidis* ; *A. pisum* and *A. gossypii*, respectively. The average number of aphids consumed per larva for the coccinellid predator was also significantly different.

The aphid species have a significant effect on the longevity of adult stage of the coccinellid predator and the fecundity of this female. The average number of eggs deposited per *H. tridecimpunctata* female was 948.29 ± 15.76 ; 815.67 ± 7.68 ; 989.54 ± 16.11 and 728.54 ± 6.85 eggs when reared on *S. avenae* ; *R. maidis*; *A. pisum* and *A. gossypii*, respectively. The statistical analysis showed that the aphid species have a highly significant effect on the female fecundity. It can be recommended that the coccinellid predator used as biological control agents against the four aphid species tested.

Keywords: Biology, *Hippodamia tridecimpunctata* L.; *Sitobion avenae* Fabr.; *Rhopalosiphum maidis* (Fitch); *Acyrtosiphon pisum* (Harris) and *Aphis gossypii* Glover.

INTRODUCTION

In recent years much interest has been given to biological control of harmful pests specially predator insects. However, the success of these biological control agents necessitates their presence in sufficient numbers when required. These obstacles can be avoided by their rearing on natural preys or artificial diets under controlled conditions for biological control of aphids and other soft-bodied insects on several economic crops (Majerus, 1994). Mass production of coccinellid predatory insects in biological control programs requires huge numbers at low costs. It is desirable to choose the

predator, which has short developmental time, a high survival rate and a high reproductive capacity.

Many authors supported the use of safe alternative control methods such as biological control in some cases. Predators as one of the major group of natural enemies play a noticeable role against different insect pests. Predators belonging to family Coccinellidae, comprise one of the most active groups of predatory species, that feed on different sucking pests including aphids, whiteflies, Jassids and mites as well as other soft bodied insect. This family gained the interest of many investigators as important group of predators in the biological control of insect pest attacking different crop plant (Hamed and Hassanein 1984; Fang *et al.* 1984; Goven and Ozgur 1990; Abdel-Samad, 1996; Abney *et al.* 2000; Samban 2003; Delrio *et al.* 2004; Hesler, *et al.* 2004 and Bahy El-Din, 2006).

The efficiency of the coccinellid predator *Hippodamia tridecimpunctata* L. as a biological control agents has been studied by few number of investigators in different parts of the world. (Radriguez and miller 1995; Hafez 2001 Mohamed 2001; and Prasifka *et al.* 2004).

Therefore, this study was undertaken to evaluate the effect prey types on certain biological characteristics of *H. tridecimpunctata* under constant temperature and relative humidity.

MATERIALS AND METHODS

Experiments were carried out at the laboratory of Economic Entomology Department, Faculty of Agriculture, Mansoura University during the period from March 2010 to May 2011 under 26 ± 1 °C and 65 ± 5.0 % R.H. . Four aphid species namely, *Sitobion avenae* Fabr. , *Rhopalosiphum maidis* (Fitch), *Acyrtosiphon pisum*, (Harris) and *Aphis gossypii* Glover were used as preys for the coccinellid predator, *H. tridecimpunctata*. The predator and the prey individuals were obtained from a maintained culture in the Insectary.

Larval experiments :

Newly hatched predator larvae from this coccinellid was put individually in a Petri-dish (10 cm. diameter) with a filter paper on its bottom. Twenty replicates from *H. tridecimpunctata* was reared on the four aphid species. Known surplus numbers of prey species were offered and the devoured individuals were replaced daily. Attacked prey individuals were counted and recorded daily throughout the periods of the larval instars. The duration period of larval and pupal stages; consumption rate and the percentage mortality were estimated.

Adult experiments :

After emergence from the pupae, the predator adults were sexed and then introduced singly into a Petri-dish. Known numbers of the four aphids were offered daily on a plant leaflet to each predator. Counting and removing the un-devoured aphids in each Petri-dish were practiced before introducing the new aphid individuals. After five days of emergence, copulation took place and the two sexes were immediately separated and kept singly in the dishes. Daily numbers of laid eggs per predator female during its ovipositional

period was counted. In addition, the total number of eggs laid per predator female was estimated. The daily averages of prey consumption throughout adults longevities were calculated.

Data analysis :

Data for the developmental time of immature stages mortality, pre-oviposition, oviposition, post-oviposition periods, fecundity, male longevity, consumption rate of the coccinellid predator reared on the tested aphid species were subjected for one way analysis of variance (ANOVA), and the means were separated using Duncan`s Multiple Range Test (CoHort Software, 2004).

RESULTS AND DISCUSSION

Reared on *Sitobion avenae*

Immature stages

Data in Table (1) indicated that the incubation period of the predator eggs lasted 3.46 ± 0.15 days. The duration period of the larval stage averaged 11.74 ± 1.46 days. The average of the total consumption during the four larval instars were 18.53 ± 1.54 , 70.12 ± 2.79 , 137.40 ± 3.85 and 469.75 ± 6.98 aphid individuals, respectively, and the average of total consumption per larva was 695.80 ± 10.75 aphid individuals. The percentages of feeding capacity for each of the four larval instars of this predator were 2.66, 10.08, 19.75 and 67.51 respectively, Therefore it may be obvious that the third and fourth instar larvae of this predator represent together 87.26 % the back one in predation activity. The percentages of mortality were 10.30, 6.52, 4.75 and 3.62 during the larval instars.

Pupal stage.

As clearly shown in Table (1), the pupal period lasted for an average of 7.46 ± 1.25 days.

Table (1): Duration period of larval and pupal stages, feeding capacity and percentage of mortality for *H. tridecimpunctata* reared on *S. avenae* under constant temperature and relative humidity.

Immature stages	Duration in days	Daily aver. consumption	Aver. of total consumption	% feeding capacity	%of mortality
A: Eggs incubation period	3.46 ± 0.15	-	-	-	-
B. Larval stage					
1 st inst.	2.36 ± 0.42	7.85 ± 0.67	18.53 ± 1.54	2.66	10.30
2 nd inst.	1.92 ± 0.36	36.52 ± 2.79	70.12 ± 2.79	10.08	6.52
3 rd inst.	2.10 ± 0.51	65.43 ± 4.56	137.40 ± 3.85	19.75	4.75
4 th inst.	5.36 ± 0.80	87.64 ± 6.25	469.75 ± 6.98	67.51	3.62
Total	11.74 ± 1.46	59.27 ± 5.37	695.80 ± 10.75	100	-
C: Pupal stage	7.46 ± 1.25	-	-	-	-

Predator adult stage

Female.

The predators female fed on a total average of 5463.59±60.47 aphid individuals, with a daily rate of 109.53±3.78 during the longevity period of 49.88±3.56 days as seen in Table (2). The average of the pre-oviposition period was 6.89±0.82 days. The predator female consumed during this period 640.80±8.29 with a daily rate of 93.00±2.76 aphid individuals. The predators female consumed during the oviposition period an average 4250.25±46.54 aphid individuals, this period lasted an average of 34.64±2.46 days with a daily rate of 122.70±4.67. The number of deposited eggs per predator female averaged 948.29±15.76 eggs with a daily rate of 27.38±2.4 per day. During the post-oviposition period the female consumed 572.54±7.95 aphid individuals, this period lasted an average of 8.35±1.10 days with a daily rate of 68.57±1.85. The results showed that the feeding capacity of predator female was always higher than that of the male as the female fed on 1.67 times of *S. avenae* more than the male (Table 2).

Male

During its longevity period which lasted for an average of 37.24±2.63 days (Table 2). The predator adult male consumed a total average 3265.47±42.56 aphid individuals with a daily rate of 87.69±2.83.

Table (2): Longevity, feeding capacity and fecundity of the predator *H. tridecimpunctata* adult reared on *S. avenae* under constant temperature and relative humidity.

Adult stage		Period in days	Daily aver. consumption	Aver. of total consumption	No. of eggs	
					Daily	Total
Female	Pre-oviposition	6.89±0.82	93.00±2.76	640.80±8.29		
	Oviposition	34.64±2.46	122.70±4.67	4250.25±46.54	27.38	948.29
	Post-oviposition	8.35±1.10	68.57±1.85	572.54±7.95	±2.4	±15.76
	Longevity	49.88±3.56	109.53±3.78	5463.59±60.47		
B. Male	Longevity	37.24±2.63	87.69±2.83	3265.47±42.56		

Reared on *Rhopalosiphum maidis*

Immature stages

Data in Table (3) showed that the incubation period of the predator eggs lasted 3.54±0.54 days. The duration period of the larval stage averaged 13.17±2.42 days. The average of the total consumption during the four larval instars were 24.92±1.30, 90.35±2.96, 146.72±5.76 and 519.82±9.46 aphid individuals, respectively, and the average of total consumption per larva was 781.83±15.68 aphid individuals. The percentages of feeding capacity for each of the four larval instars of this predator were 3.19, 11.56, 18.77 and 66.48 respectively, Therefore it may be obvious that the third and fourth instar larvae of this predator represent together 85.25 % the back one in predation activity. The percentages of mortality were 10.62, 6.81, 5.10 and 4.27 during the larval instars.

Pupal stage.

As clearly shown in Table (3), the pupal period lasted for an average of 7.83±1.32 days.

Table (3): Duration period of larval and pupal stages, feeding capacity and percentage of mortality for *H. tridecimpunctata* reared on *R. maidis* under constant temperature and relative humidity.

Immature stages	Duration in days	Daily aver. consumption	Aver. of total consumption	% feeding capacity	%of mortality
Eggs incubation period	3.54±0.54	-	-	-	-
B. Larval stage					
1 st inst.	2.54±0.54	9.82±0.94	24.92±1.30	3.19	10.62
2 nd inst.	2.16±0.92	41.83±2.67	90.35±2.96	11.56	6.81
3 rd inst.	2.37±0.85	61.91±4.52	146.72±5.76	18.77	5.10
4 th inst.	6.10±1.26	85.22±3.42	519.82±9.46	66.48	4.27
Total	13.17±2.42	59.36±4.12	781.83±15.68	100	-
C:Pupal stage	7.83±1.32	-	-	-	-

Predator adult stage

Female.

The predator female fed on a total average of 6056.06±100.65 aphid individuals, with a daily rate of 129.54±5.62 during the longevity period of 46.75±3.78 days as seen in Table (4). The average of the pre-oviposition period was 7.12±0.80 days. The predator female consumed during this period 694.74±8.56 with a daily rate of 97.58±4.67 aphid individuals. The predators female consumed during the oviposition period an average 4764.62±80.67 aphid individuals, this period lasted an average of 32.78±2.67days with a daily rat of 145.35±6.45. The number of deposited eggs per predator female averaged 815.67±7.64 eggs with a daily rate of 24.88±1.96 per day. During the post-oviposition period the female consumed 596.70±7.35 aphid individuals, this period lasted an average of 6.85±0.56 days with a daily rate of 87.11±3.62. The results showed that the feeding capacity of predator female was always higher than that of the male as the female fed on 1.75 times of *R. maidis* more than the male (Table 4).

Male

During its longevity period which lasted for an average of 34.24±2.96 days (Table 4). The predator adult male consumed a total average 3457.88±70.68 aphid individuals with a daily rate of 100.99±4.22.

Table (4): Longevity, feeding capacity and fecundity of the predator *H. tridecimpunctata* adult reared on *R. maidis* under constant temperature and relative humidity.

Adult stage		Period in days	Daily average consumption	Aver. total consumption	No. of eggs	
					Daily	Total
Female	Pre-oviposition	7.12±0.80	97.58±4.67	694.74±8.56		
	Oviposition	32.78±2.67	145.35±6.45	4764.62±80.67	24.88	815.67
	Post-oviposition	6.85±0.56	87.11±3.62	596.70±7.35	±1.96	±7.64
	Longevity	46.75±3.78	129.54±5.62	6056.06±100.65		
Male	Longevity	34.24±2.96	100.99±4.22	3457.88±70.68		

Reared on *Acyrtosiphon pisum*

Immature stages

Data in Table (5) indicated that the incubation period of the predator eggs lasted 3.26±0.10 days. The duration period of the larval stage averaged

10.44±1.15 days. The average of the total consumption during the four larval instars were 16.85±1.42, 64.44±3.56, 120.68±5.72 and 410.53±7.56 aphid individuals, respectively, and the average of total consumption per larva was 612.50±8.96 aphid individuals. The percentages of feeding capacity for each of the four larval instars of this predator were 2.75, 10.52, 19.70 and 67.03 respectively, Therefore it may be obvious that the third and fourth instar larvae of this predator represent together 86.73 % the backbone in predation activity. The percentages of mortality were 9.32, 4.64, 3.51 and 2.45 during the larval instars.

Pupal stage.

As clearly shown in Table (5), the pupal period lasted for an average of 6.97±1.21 days.

Table (5): Duration period of larval and pupal stages, feeding capacity and percentage of mortality for *H. tridecimpunctata* reared on *A. pisum* under constant temperature and relative humidity.

Immature stages	Duration in days	Daily aver. consumption	Ave. of total consumption	% feeding capacity	% of mortality
Eggs incubation period	3.26±0.10	-	-	-	-
Larval stage					
1 st inst.	2.08±0.60	8.10±0.97	16.85±1.42	2.75	9.32
2 nd inst.	1.54±0.23	41.84±2.67	64.44±3.56	10.52	4.64
3 rd inst.	1.97±0.35	61.26±4.25	120.68±5.72	19.70	3.51
4 th inst.	4.85±0.90	84.65±5.62	410.53±7.56	67.03	2.45
Total	10.44±1.15	58.67±3.26	612.50±8.96	100	-
C:Pupal stage	6.97±1.21	-	-	-	-

Predator adult stage

Female.

The predator female fed on a total average of 4933.73±25.67 aphid individuals, with a daily rate of 96.17±3.62 during the longevity period of 51.30±2.96 days as seen in Table (6). The average of the pre-oviposition period was 6.53±0.86 days. The predator female consumed during this period 587.34±3.56 with a daily rate of 89.94±1.75 aphid individuals. The predator female consumed during the oviposition period an average 3925.74±18.64 aphid individuals, this period lasted an average of 35.67±1.78 days with a daily rate of 110.06±2.89. The number of deposited eggs per predator female averaged 989.54±12.57 eggs with a daily rate of 27.74±2.12 per day. During the post-oviposition period the female consumed 420.65±2.75 aphid individuals, this period lasted an average of 9.10±0.97 days with a daily rate of 46.26±1.25. The results showed that the feeding capacity of predator female was always higher than that of the male as the female fed on 1.72 times of *A. pisum* more than the male (Table 6).

Male

During its longevity period which lasted for an average of 36.27±2.10 days (Table 6). The predator adult male consumed a total average 2874.76±18.11 aphid individuals with a daily rate of 79.26±2.75.

Table (6): Longevity, feeding capacity and fecundity of the predator *H. tridecimpunctata* adult reared on *A. pisum* under constant temperature and relative humidity.

Adult stage		Period in days	Daily aver. consumption	Aver. of total consumption	No. of eggs	
					Daily	Total
Female	Pre-oviposition	6.53±0.86	89.94±1.75	587.34±3.56		
	Oviposition	35.67±1.78	110.06±2.89	3925.74±18.64	27.74	989.54
	Post-oviposition	9.10±0.97	46.26±1.25	420.65±2.75	±2.12	±12.57
	Longevity	51.30±2.96	96.17±3.62	4933.73±25.67		
Male	Longevity	36.27±2.10	79.26±2.75	2874.76±18.11		

**Reared on *Aphis gossypii*
Immature stages**

Data in Table (7) showed that the incubation period of the predator eggs lasted 3.35±0.12 days. The duration period of the larval stage averaged 13.81±1.84 days. The average of the total consumption during the four larval instars were 27.84±1.76, 89.72±2.85, 180.35±4.26 and 596.62±7.52 aphid individuals, respectively, and the average of total consumption per larva was 894.53±11.61 aphid individuals. The percentages of feeding capacity for each of the four larval instars of this predator were 3.11, 10.03, 20.16 and 66.70 respectively, Therefore it may be obvious that the third and fourth instar larvae of this predator represent together 86.86 % the back one in predation activity. The percentages of mortality were 11.96, 7.64, 5.87 and 4.65 during the larval instars.

Pupal stage.

As clearly shown in Table (7), the pupal period lasted for an average of 8.26±1.42 days.

Table (7): Duration period of larval and pupal stages, feeding capacity and percentage of mortality for *H. tridecimpunctata* reared on *A. gossypii* Glover under constant temperature and relative humidity.

Immature stages	Duration in days	Daily aver. consumption	Aver. of total consumption	% feeding capacity	% of mortality
Eggs incubation period	3.35±0.12	-	-	-	-
Larval stage					
1 st inst.	2.94±0.65	9.47±0.96	27.84±1.76	3.11	11.96
2 nd inst.	2.10±0.49	42.72±2.54	89.72±2.85	10.03	7.64
3 rd inst.	2.35±0.54	76.74±3.76	180.35±4.26	20.16	5.87
4 th inst.	6.42±0.96	92.93±5.35	596.62±7.52	66.70	4.65
Total	13.81±1.84	64.77±3.16	894.53±11.61	100	-
C:Pupal stage	8.26±1.42	-	-	-	-

**Predator adult stage
Female.**

The predator female fed on a total average of 6608.86±48.56 aphid individuals, with a daily rate of 125.79±3.84 during the longevity period of 52.54±4.09 days as seen in Table (8). The average of the pre-oviposition period was 7.32±0.96 days. The predator female consumed during this period 819.42±5.78 with a daily rate of 111.94±3.58 aphid individuals. The predators

female consumed during the oviposition period an average 5142.17±26.56 aphid individuals, this period lasted an average of 36.26±3.11 days with a daily rate of 141.81±7.67. The number of deposited eggs per predator female averaged 728.54±6.85 eggs with a daily rate of 20.09±1.56 per day. During the post-oviposition period the female consumed 647.27±4.80 aphid individuals, this period lasted an average of 8.96±0.45 days with a daily rate of 72.24±2.50. The results showed that the feeding capacity of predator female was always higher than that of the male as the female fed on 1.67 times of *A. gossypii* more than the male (Table 8).

Male

During its longevity period which lasted for an average of 38.72±2.96 days (Table 8). The predator adult male consumed a total average 3948.12±32.67 aphid individuals with a daily rate of 86.47±2.85.

Table (8): Longevity, feeding capacity and fecundity of the predator *H. tridecimpunctata* adult reared on *A. gossypii* Glover under constant temperature and relative humidity.

Adult stage		Period in days	Daily aver. consumption	Aver. of total consumption	No. of eggs	
					Daily	Total
Female	Pre-oviposition	7.32±0.96	111.94±3.58	819.42±5.78		
	Oviposition	36.26±3.11	141.81±7.67	5142.17±26.56	20.09±	728.54
	Post-oviposition	8.96±0.45	72.24±2.50	647.27±4.80	1.56	±6.85
	Longevity	52.54±4.09	125.79±3.84	6608.86±48.56		
Male	Longevity	38.72±2.96	86.47±2.85	3948.12±32.67		

Effect of prey types on the developmental time of immature stage:

Data in (Table 9 and Figure 1) indicated that the incubation period of *H. tridecimpunctata* varied from 3.26±0.10 to 3.54± 0.54 days with significant difference among the four tested aphid species. Considering the developmental time of larval instars, the shortest developmental time was obtained when larvae reared on *A. pisum* (10.44±1.15 days), while the longest time was recorded on *A. gossypii* (13.84±1.84 days). The developmental of larval instars showed a significant variation among the four aphid species (Table 9 and Figure 1). Among the tested aphid species, there were significant differences between developmental times of pupal stage. The longest time was observed with *A. gossypii*, while the shortest time was obtained with *A. pisum*. The total developmental time (from egg hatching to adult eclosion) was ranged from 20.67±2.10 days by rearing on *A. pisum* to 25.42±3.25 days by feeding on *A. gossypii* with significantly differed. Mortality percentage from egg to adult ranged from 4.98 % when reared on *A. pisum* to 9.05 % with *S. avenae*. There were significant variations among the four tested aphid species. These findings agree with that of Lotfalizadeh *et al.*, (2000) who found that the larval stage period was 20.97 days and the pupal period lasted 6.13 days when reared on *P. vovae*. Bahy El-Din (2006) found that, the total larval durations *H. convergens* were 12.91; 12.04; 10.78 and 8.84 days when fed on 1st; 2nd; 3rd and 4th nymphal instars of *A. craccivora*. The respective total developmental period was 21.55; 20.8; 19.49 and 17.56 days

Table (9): Developmental time of the immature stages of *H. tridecimpunctata* predator reared on certain aphid species under constant temperature and relative humidity.

Prey species	Incubation period	Larval instars					Pupal stage	Total of immature stages	Mortality %
		1 st	2 nd	3 rd	4 th	Total			
<i>S. avenae</i>	3.46± 0.15a	2.36± 0.42b	1.9± 0.36a	2.1± 0.51a	5.36± 0.80b	11.74± 1.46c	7.4± 1.25b	22.66± 2.96c	9.05a
<i>R. maidis</i>	3.54± 0.54a	2.54± 0.92a	2.1± 0.82a	2.3± 0.85a	6.10± 1.26a	13.17± 2.42b	7.8± 1.32b	24.54± 2.98b	6.70c
<i>A. pisum</i>	3.26± 0.10b	2.08± 0.60c	1.5± 0.23b	1.9± 0.35a	4.85± 0.90c	10.44± 1.15d	6.9± 1.21c	20.67± 2.10d	4.98d
<i>A. gossypii</i>	3.35± 0.12b	2.94± 0.65a	2.1± 0.49a	2.3± 0.54a	6.42± 0.96a	13.84±1. 84a	8.2± 1.42a	25.42± 3.25a	7.53b

^a Means followed by the same letter in a column for each coccinellid predator are not significantly different at the 1 % level of probability (Duncan's Multiple Range Test).

Feeding capacity:

Data in (Table 10 and Figure 1) showed the consumption period rate of *H. tridecimpunctata* larval instars when reared on four aphid species. The average number of aphids consumed during first instar larvae varied from 16.85±1.42 individuals of *A. pisum* to 27.84±1.76 individuals of *A. gossypii*. Consumption percentage differed between 2.66 % on *S. avenae* to 3.19 % on *R. maidis*. There were apparent differences in average numbers of the four consumed aphid species and consumption percentage by the second instar larvae. In addition, the same trend was obtained with the third and fourth instars larvae. The total consumption rate from the four aphid species by the larval stage of *H. tridecimpunctata* showed significant difference. Concerning to the food preference for predator larvae among tested aphid species, *A. gossypii* came first, followed by *R. maidis*, *S. avenae* and *A. pisum*. The average number of consumed aphid per larvae was also significantly different.

Table (10): Mean numbers consumed and percentage from different aphid species during larval instars of *H. tridecimpunctata* under constant temperature and relative humidity.

Prey species	Larval instars								Total
	1 st		2 nd		3 rd		4 th		
	No.	%	No.	%	No.	%	No.	%	
<i>S. avenae</i>	18.53± 1.54c	2.66b	70.12± 2.79b	10.08b	137.40± 3.85c	19.75b	469.75± 6.98c	67.51a	695.80± 10.75c
<i>R. maidis</i>	24.92± 1.30b	3.19a	90.35± 2.96a	11.56a	146.72± 5.76b	18.77c	519.82± 9.46b	66.48b	781.83± 15.68b
<i>A. pisum</i>	16.85± 1.42d	2.75b	64.44± 3.56c	10.52b	120.68± 5.72d	19.70b	410.53± 7.56d	67.03a	612.50± 9.55d
<i>A. gossypii</i>	27.84± 1.76a	3.11a	89.72± 2.85a	10.03b	180.35± 4.26a	20.16a	596.62± 7.52a	66.70b	894.53± 11.61a

^a Means followed by the same letter in a column for each coccinellid predator are not significantly different at the 1 % level of probability (Duncan's Multiple Range Test).

Longevity and fecundity of adult stage:


Data in (Table 11 and Figure 2) showed that the mean male longevity of this predator was significantly longer when fed on *A. gossypii*, than when

reared on *A. pisum*, and shortest on *S. avenae* and *R. maidis*. Concerning the female, there were significant differences in the total longevity periods of *H. tridecimpunctata* when reared on the tested aphid species (Table 11). Also, the aphid species have a significant effect on female fecundity. The highest numbers of eggs were obtained when females fed on *A. pisum* followed by *S. avenae* and *R. maidis*, while the lowest numbers were achieved on *A. gossypii* (Table 11 and Figure 2). Remzi and Bora (2002), indicated that the total number of eggs per female was 428.5 which lasted 75.3 days. Whereas Hafez, (2001) found that daily rate of *S. graminum* consumption by adult of *H. tridecimpunctata* was 34.24 individuals for female at 22.5°C. Mohamed (2001) found that *H. tridecimpunctata* fed on *A. craccivora* female consumed 108.46;122.57 and 81.56 preys during pre-ovipositional; ovipositional and post-ovipositional periods respectively. Saleh (2005) recorded that the number of eggs were 310.75 ± 11.64 eggs during the oviposition period which lasted 32.85 ± 2.97 days. El-Heneidy et al. (2008) recorded that the total life time fecundity of 880 eggs/ female for *H. convergens* when fed on *A. craccivora*. Mandoure (2009) found that *H. pruni* was suitable for development and reproduction of *H. convergens*. Abd El-Motaal(2011) reported that the total fecundity of *H. convergens* females were 1030.10; 1063.30 and 1010.20 eggs when reared on *A. craccivora*; *H. pruni* and *R. padi*.


Table (11): Longevity, feeding capacity and fecundity of *H. tridecimpunctata* reared on four aphid species under controlled conditions and relative humidity.

Aphid species	Female longevity				Male longevity	Ave. of total consumption		Female fecundity	
	Pre-ovip.	Oviposition	Post-ovip.	Total		Female	Male	Daily	Total
<i>S. avenae</i>	6.89± 0.82b	34.64± 2.46c	8.35± 1.10c	49.88 ±3.56b	34.24± 2.63c	5463.59± 60.47c	3265.47± 42.56c	27.38 ±2.4b	948.29 ±15.76b
<i>R. maidis</i>	7.12± 0.80a	32.78± 2.67d	6.85± 0.56d	46.75 ±3.78c	34.24± 2.96c	6056.06± 100.65b	3457.88± 70.68b	24.88 ±1.96c	815.67 ±7.68c
<i>A. pisum</i>	6.53± 0.86b	35.67± 1.78b	9.10± 0.97a	51.3± 2.96ab	36.27± 2.10b	4933.73± 25.67d	2874.76± 18.11d	27.74 ±2.12a	989.54 ±16.11a
<i>A. gossypii</i>	7.32± 0.96a	36.26± 3.11a	8.96± 0.45b	52.54 ±4.09a	38.72± 2.96a	6608.86± 48.56a	3948.12± 32.67a	20.09±1. 56d	728.54 ±6.85d

^a Means followed by the same letter in a column for each coccinellid predator are not significantly different at the 1 % level of probability (Duncan's Multiple Range Test).



Figure(1): Effect of prey types on certain biological characteristics of *H. tridecimpunctata* immature stages under constant temperature and relative humidity.



Figure(2): Effect of prey types on certain biological characteristics of *H. tridecimpunctata* adults under constant temperature and relative humidity.

REFERENCES

- Abd El-Motaal; D. S. M. 2011. Biological and Ecological studies on Hippodamid convergens(Coleoptra: Coccinellidae) M. Sc. Thesis; Faculty of Agriculture, Suez Canal University pp159.
- Abdel-Samad, S. M. S. 1996. Studies on natural enemies of certain insects attacking leguminous crops.M. Sc. Thesis, Fac. Of Agric., Ain Shamas Univ. 94pp.
- Abney, M. R. ; J. R. Ruberson; G. A. Herzog; T. J. Kring; D. C. Steinkraus; P. Dugger (ed.) and D. Richter 2000. Natural enemies and population dynamic of cotton aphid, *Aphis gossypii* in Georgia. Proceeding Beltwide cotton conference, San Antonio, U. S. A., 4-8 January, (2): 1246-1247.
- Bahy El-Din, I. A. 2006. Studies on the biology and feeding capacity of some coccinellids species. M. Sc.Thesis. Fac. Agric. Moshtohor, Benha Univ. 158 pp.
- CoHort Software, 2004. CoStat. www. CoHort com. Monterey, California, U. S. A.
- Delrio, G.; M. verdinelli and G. Serra, 2004. Monitoring of pest and beneficial insect populations in summer sown *Bt* maize. Bulletin-OILB/SROP. 27(3):43-48.
- El-Heneidy, A. H.; A. A. Hafez; F. F. Shalaby and I. A. B. Bahy-Eldin, 2008. Comparative biological aspects of two coccinellid species; *Coccinella undecimpunctata* L. and *Hippodamia convergens* Guer. Under laboratory conditions. Egyptian Journal of Biological Pest Control, 18(1):51-59.
- Fang, C. Y. ; S. G. Wen; S. Z. Cui and Y. H. Wang, 1984. The role of natural enemies in the integrated control of insect pests on cotton. China Cotton. (2): 42-43.
- Goven, M. A. and A. F. Ozgur, 1990. The influence of natural enemies on populations of *Thrips tabaci* on cotton in the southeast region of Anatolia. Proceedings of the Second Turkisk National Congress of Biological Control, 155-164.

- Hafez, A. A. 2001. Functional response, and some biological parameters of *Hippodamia tridecimpunctata* Goet. (Coleoptera: Coccinellidae). Egypt. J. Biol. Pest Control 11(1): 31-38.
- Hamed, A. R. and F. A. Hassanein, 1984. Assessment of the role of *Coccinella undecimpunctata* L. (Col., Coccinellidae) as a biological control agent against *Spodoptera littoralis* Boisd. Z. Angew. Entomol. 97(5): 520-523.
- Hesler, L. S.; R. W. Kieckhefer and A. M. Catangui, 2004. Surveys and field observations of *Harmonia axyridis* and other coccinellidae (Coleoptera) in Eastern and central south Dakota. Trans. Am. Entomol. Soc., 130(1): 133-133.
- Lotfalizadeh, H.; B. Hatami and J. Khalaghami 2000. Biology of *Exochomus quadripustulatus* (L) (Coleoptera: Coccinellidae) on Cypress tree mealy bugs *Planococcus vovae* (Nasanov) (Homoptera: Pseudococcidae) in Shiraz. J. Entomol. Soci. Iran, 20 (1): 61-76.
- Majerus M. E. N. 1994. Lady birds. Harper Collins, London.
- Mandour, N. S. 2009. Development, survival, feeding consumption and reproduction of the convergent lady beetle, *Hippodamia convergens* (Coleoptera: Coccinellidae) feeding on three aphid species. Agricultural Research Journal, Suez Canal University, 9(1):103-108.
- Mohamed, N. A. 2001. Mass-rearing of certain predatory insects on artificial diets for controlling some insects infesting vegetable crops. Ph. D. Thesis, Fac. Agric., Mansoura Univ., 93 pp.
- Prasifka, J. R.; K. M. Heinz and K. O. Winemiller, 2004. Crop colonization, feeding and reproduction by the predatory beetle, *Hippodamia convergens* as indicated by stable carbon isotope analysis. Ecol. Entomol. 29(2): 226-233.
- Remzi, A. M. and K. Bora, 2002. Development, survival and reproduction of three coccinellids feeding on *Hyalopterus pruni* (Geoffe) (Homoptera: Aphididae). Turk J. Agric. 26: 119 – 124 .
- Rodriguez, S. C. and J. C. Miller, 1995. Life history traits in *Hippodamia convergens* (Coleoptera: Coccinellidae) after selection for fast development. Biol. Control, 5(3): 389-396.
- Saleh, , M. F. 2005. Biological studies of certain predatory insects at Mansoura district. M. Sc. Thesis. Fac. Agric. Mansoura Univ. pp. 149.
- Samban, H. M. I. 2003. The population fluctuations of certain predaceous arthropods inhabiting three field crops in Minia. Assiut J. Agrig. Sci. 34(2):183-191.

تأثير أنواع الفرائس على بعض الخصائص البيولوجية لأبو العيد ذو ثلاثة عشر
نقطة *Hippodamia tridecimpunctata* L. تحت درجة حرارة و رطوبة
ثابتة

نادية الحسيني محمد^١ و عبد البديع عبد الحميد غانم^٢ و أحمد أمين أحمد صالح^٢
^١ معهد بحوث وقاية النباتات - مركز البحوث الزراعية - الدقي - الجيزة - مصر .
^٢ قسم الحشرات الاقتصادية - كلية الزراعة - جامعة المنصورة .

اجريت تجارب معملية لدراسة بعض الخصائص البيولوجية لمفترس أبو العيد ذو ثلاثة عشر نقطة *H. tridecimpunctata* عند تربيته على أنواع المن التالية من الحبوب *Sitobion avenae* Fabr و من الذرة *Rhopalosiphum maidis* (Fitch) و من البسلة *Acyrtosiphon pisum* (Harris) و من القطن *Aphis gossypii* Glover تحت درجة حرارة و رطوبة ثابتة و كانت النتائج المتحصل عليها توضح مدة التطور من فقس البيض حتى خروج الحشرات الكاملة لهذا المفترس أظهرت تأثيرا معنويا واضحا عند التربية على أنواع المن السابقة كما أن النسبة المئوية للموت لهذا المفترس من فقس البيض و حتى خروج الحشرة الكاملة تراوحت بين ٤.٨٩% عند التغذية على من البسلة إلى ٩.٠٥% عند التغذية على من الحبوب و أوضح التحليل الإحصائي اختلافات نسبة الموت معنويا بين أنواع المن الأربعة و أكدت النتائج أن معدل الإستهلاك الكلي لكل يرقة من المفترس لأنواع المن الأربعة كانت كالآتي ١٠.٧٥±٦٩٥.٨ و ١٠.٨٣±٧٨١.٨٣ و ١٠.٧٥±٦١٢.٥ و ٩.٥٥±٨٩٤.٥٣ فرد من المن عند التربية على من الحبوب و من الذرة و من البسلة و من القطن على التوالي و كان متوسط عدد المن المستهلك لكل يرقة لهذا المفترس مختلف معنويا لأنواع المن الأربعة السابقة كما أثرت أنواع المن المختلفة تأثيرا معنويا على فترة حياة الطور الكامل لهذا المفترس كذلك الكفاءة التناسلية لإناثه فكان متوسط عدد البيض الذي تم وضعه للإناث ١٥.٧٦±٩٤٨.٢٩ و ٧.٧٨±٨١٥.٦٧ و ١٦.١١±٩٨٩.٥٤ و ٦.٨٥±٧٢٨.٥٤ بيضة عند التربية على من الحبوب و من الذرة و من البسلة و من القطن على التوالي و أكد التحليل الإحصائي أن أنواع المن الأربعة التي تم دراستها كان تأثيرها على المعنوية على الكفاءة التناسلية للإناث و توضح النتائج أن من البسلة كان أفضل أنواع المن المختبرة لهذا المفترس حيث وضعن إناثه أكبر كمية من البيض عليه لذلك ننصح بأن مفترس أبو العيد ذو ثلاثة عشر نقطة *H. tridecimpunctata* يمكن استخدامه كعنصر من عناصر مكافحة البيولوجية لأنواع المن الأربعة

قام بتحكيم البحث

كلية الزراعة - جامعة المنصورة
كلية الزراعة - جامعة القاهرة

أ.د / هاله احمد كامل الصيرفي
أ.د / محمود السيد نور

