

- A. Diabetes insipidus
- B. Seizures
- C. Papilledema
- D. Facial weakness

124. Fibrous dysplasia is associated with what syndrome?

- A. Turcot's syndrome
- B. Gardner's syndrome
- C. McCune-Albright syndrome
- D. Paget's disease

125. You are operating on a cerebellar hemangioblastoma with a large associated cystic component. You open the dura and the cerebellum begins to herniate through the dural defect. What will be the most effective means to decrease posterior fossa pressure?

- A. Hyperventilation
- B. Mannitol
- C. Dexamethasone
- D. Needle aspiration of cystic contents

126. You resect a pathology proven cerebellar hemangioblastoma with a large cystic component. You have removed the mural nodule. Should you attempt to excise the entire cyst wall?

- A. Yes
- B. No

127. You resect a pathology proven cerebellar pilocytic astrocytoma with a large cystic component. You have removed the mural nodule. Should you attempt to excise the entire cyst wall?

- A. Yes
- B. No

128. During endoscopic third ventriculostomy, aggressive manipulation of the endoscope within the third ventricle should be avoided to prevent injury to what structure?

- A. Mamillary bodies
- B. Caudate head
- C. Fornix
- D. Thalamus

129. Approximately what length of temporal lobe can be resected safely during a temporal lobectomy on the dominant side?

- A. 1 to 2.5 cm
- B. 3 to 4.5 cm