

## The Role of Sulphur Application and Dicyandiamide Comparing with Mineral Fertilization Npk on Chemical Constituent of Lettuce Plant (*Lactuca Sativa L.*)

Fouda, K. F.

Soil Sci. Dept., Fac. of Agric., Mansoura Univ., Egypt.



### ABSTRACT

A field experiment was conducted in a clayey soil at the experimental farm of the Faculty of Agric. El-Mansoura Univ. to study the effect of sulphur (S) application and Dicyandiamide, (DCD) comparing with mineral fertilization of NPK rates on chemical composition of lettuce plant (*Lactuca sativa L.*) in factorial design. Experiment included 9 treatments which were 3 rates of NPK fertilization (50, 75 and 100% from recommended dose) and 3 treatments of (DCD, S and DCD + S) with 3 replicates. Thus, the total number of the experimental plots was 27 plots. Results indicated that chlorophyll content, N, P, K and S concentrations,  $\text{NO}_3\text{-N}$  and  $\text{NO}_2\text{-N}$  content  $\text{mg.l}^{-1}$  as well as available N, P and K  $\text{mg.l}^{-1}$  in leaves were significantly increased due to increasing NPK fertilization rates, and the highest mean values were recorded with adding 100% NPK from recommended dose, while nitrate reductase activity was decreased. The highest average values of mentioned parameters indicated that using mix of (S & DCD) comparing with other treatments except  $\text{NO}_3\text{-N}$ , and  $\text{NO}_2\text{-N}$   $\text{mg.l}^{-1}$  which recorded with adding sulphur. In the same time the effect of interaction among treatments on the traits illustrated that the highest values of all parameters in lettuce were connected with 100% NPK + S + DCD. While, nitrate reductase activity of lettuce recorded the highest mean values with S+DCD+ 50% NPK over the control

### INTRODUCTION

Lettuce (*Lactuca sativa L.*) belongs to the botanical family Asteraceae and is an option in the olericulture for the cultivation both in protected environment and at the field. Commonly, the produced leaves are commercialized and freshly consumed in the form of salad, which is an alternative of healthy food in the Brazilian menu (Midan and Sorial, 2011). Lettuce leaves are a rich source of antioxidants, vitamins A and C, and phytochemicals which are anti-carcinogenic (Masarirambi *et al.*, 2012). The cultivated area of lettuce in Egypt is about 3110 hectares, which produced about 68644 tons (MALR, 2012).

Mineral fertilizers were used to provide soil nutrients in order to maintain optimum soil fertility conditions and healthy growth of plants and quality yield. NPK fertilizers are required greatly by crops for healthy development and crop quality. Furthermore, N is an important factor for higher yield and average head weight of lettuce (Hosseney and Ahmed, 2009). However, the increase in the nitrogen fertilization rate enables obtaining a higher yield but at the same time conveys the risk of deteriorating yield quality resulting from an excessive nitrate accumulation. It particularly refers to leaf vegetables. Phosphorus (P) is the second major nutrient for plant growth as it is an integral part of different biochemical like nucleic acids, nucleotides, phospholipids and phosphoproteins. Inadequate P supply will result in a decreased synthesis of RNA, the protein maker, leading to decreased growth. Grain yield is often severally reduced with P deficiency (Jones *et al.*, 2003). Potassium is required in least amount but in soil it is required in large amount by many crops and it is important for maintaining the osmotic potential and rigidity of plant cells; hence it plays a vital role in water relations in the plant (Okoli and Nweke, 2015).

Dicyandiamide (DCD containing 66.7% N) is the most widely used nitrification inhibitor because it is cheaper, nontoxic, less volatile, relatively water soluble, and relatively benign to non-target microbial

communities (O'Callaghan *et al.*, 2010 and Di *et al.*, 2011). Many studies have shown that DCD can significantly decrease  $\text{NO}_3$  leaching and  $\text{NO}_2$  emissions from cropping systems or grazed pasture systems (Di *et al.*, 2007 and Cui *et al.*, 2011). The nitrification inhibitor DCD decreases  $\text{NO}_3^-$  leaching by inhibiting the growth and activity of the ammonia-oxidizing bacteria in the soil, thus slowing down the rate of nitrification and keeping the N in the  $\text{NH}_4^+$  form which is adsorbed onto the soil exchange surfaces and is available for plant uptake (Asing *et al.*, 2008, Di *et al.*, 2010).

Sulfur (S) is one of nature's super nutrients and one of the oldest elements known. It is the thirteenth most abundant element in the earth's crust. S is essential for the production of three amino acids found in both plants and animals. These amino acids are the building blocks in the synthesis of proteins. Without proteins, plants and animals simply could not exist. Mehana and Farag (2000) and El Sayed *et al.*, (2015) mentioned that adding elemental sulphur had an important role in soil pH reduction, that increased availability of some nutrient elements such as P, Fe, Zn, Mn and Cu which were reflected on plant uptake and plant growth.

The main objective of the present investigation was to study the effect of sulphur (S) application and Dicyandiamide, (DCD) comparing with mineral fertilization NPK rates on chemical constituent of lettuce plant (*Lactuca sativa L.*)

**Keywords:** sulphur, Dicyandiamide, NPK fertilization and lettuce plant

### MATERIALS AND METHODS

A field experiment was conducted in a clayey soil at the experimental farm of the Faculty of Agric. El-Mansoura Univ. to investigate the application effect of sulphur (S) and Dicyandiamide, (DCD) comparing with mineral fertilization with NPK on chemical composition of lettuce plant (*Lactuca sativa L.*).

The soil of the experimental field was clayey with pH 7.9, EC:  $3.87 \text{ dSm}^{-1}$ ,  $\text{CaCO}_3$ :  $36.8 \text{ g kg}^{-1}$ , OM:  $17.6 \text{ g kg}^{-1}$ . Available N, P and K contents were 43.4,

5.21 and 190 mg kg<sup>-1</sup> during the season of the experiment, respectively which determined according to Sahlemedhin and Taye (2000). Experiment included 9 treatments which were 3 rates of NPK fertilization (50, 75 and 100%) and 3 treatments of (DCD, S and DCD + S) with 3 replicates. Thus, the total number of the experimental plot was 27 plot comparing with 100% NPK from recommended doses as control.

Seeds of lettuce cv. dark green, were sown on January 5<sup>rd</sup>, at 30 cm apart between each seedling and the other on both sides of ridgs (3m long and 70 cm wide). Each plot consisted of 3 ridgs making an area of about 6.3 m<sup>2</sup>.

Mineral fertilizers used in this investigation were urea (U) (46.5% N), calcium super phosphate (7 % P) and potassium sulphate (40 % K). N fertilizer was added in 2 equal doses directly before the first and second irrigation, while the P fertilizer was added during soil preparation for sowing and K fertilizer was given before the first irrigation. Dicyandiamide 4% N was added as a nitrification inhibitor by mixing it with nitrogen fertilization (U). Sulphur rate was added at the rate of 200 kg fed<sup>-1</sup> sulphur twice after 20 days from sowing and a week after.

At marketing stage; 90 days after sowing (May 3<sup>rd</sup>) during the season of lettuce seedling, five plants were randomly taken from each plot. The plants were weighed and then determined nutrient, where N, P, K, S% (Mertens, 2005a & b), NO<sub>3</sub>-N, NO<sub>2</sub>-N mg.l<sup>-1</sup> and nitrate reductases activity (Singh, 1988 and Hageman and Reed, 1980) as well as chlorophyll a and b (Gavrilenko and Zigalova (2003), then calculate the total chlorophyll. Soil available forms of N, P and K mg.l<sup>-1</sup> were measured in the soil samples after harvesting as determined by Haluschak, 2006 and Reeuwijk, (2002).

Data reported for chemical content of lettuce plant and soil availability were assessed by analyses of variance (ANOVA), Duncan's and the least significant difference (L.S.D) method was used for any significant differences at the P< 0.05 levels between the means of treatment values to the methods described by Gomez and Gomez, (1984). All the analyses were conducted using software computer CoSTATE.

## RESULTS AND DISCUSSION

### • Chlorophyll contents:

Results presented in Table 1 declare the effect of various concentrations of NPK fertilization, sulphur and DCD and their interactions comparing with treatment of 100% NPK from recommended dose on photosynthetic pigments during 2016.

Data showed generally, that there was an increasing in chlorophyll a, b, and a+b with increasing NPK fertilization. The increasing of change in pigments was significantly increased by increasing NPK levels. The significant increases were found in plants grown under high NPK levels i.e. 0.668, 0.481 and 1.149 mg g<sup>-1</sup> FW of Chl a, Chl b and total, respectively over rate of 100% NPK from recommended dose. The satisfactory effects of mineral application on the content

of chlorophyll may be attributed to its prominent role in the chlorophyll pigment synthesis or chlorophyll molecule in the plant tissues. These findings are corresponding to those reported by Saif-El-Deen (2000); Nabih (2002) and Abd-Elfattah, (2012).

In the same table using individual rates of DCD, S and S + DCD significantly affected chlorophyll a, b, and a+b, the highest average values were indicated with using mix of S + DCD comparing with other treatments. These results may be due to the beneficial effect of the applied-S as a one of many elements required for plant growth, its important in the formation of protein and chlorophyll (Morris *et al.*, 1984). The obtained results are in harmony with those reported by Cuppett *et al.*, (1999), Al-Redhaiman *et al.*, (2003); Wright *et al.*, (2010) and Sanchez and Silvertooth (2016).

It is obvious from the same data in Table 1 that all characteristics are affected by interaction. In general, plants received sulphur at 200 kg fed<sup>-1</sup> with DCD gave the highest values of chlorophyll content investigated during the season under all treatments of NPK fertilization comparing with 100% NPK from recommended dose (control). The highest values of traits were recorded with 100% NPK+ S + DCD.

**Table1. Effect of sulphur application, Dicyandiamide (DCD) and NPK mineral fertilization rates on chlorophyll content.**

Char. Treat.	Chlorophyll a mg g <sup>-1</sup> FW	Chlorophyll b mg g <sup>-1</sup> FW	Total a+b chlorophyll mg g <sup>-1</sup> FW
NPK fertilization			
50% RD	0.638c	0.450c	1.088c
75% RD	0.659b	0.470b	1.129b
100% RD	0.668a	0.481a	1.149a
LSD <sub>at 5%</sub>	0.002	0.006	0.006
S, DCD and S+DCD fertilization			
DCD	0.650b	0.461b	1.111b
S	0.629c	0.442c	1.071c
DCD +S	0.686a	0.497a	1.183a
LSD <sub>at 5%</sub>	0.005	0.005	0.007
Interaction			
100% RD	0.650f	0.460f	1.110f
50% DCD	0.623i	0.434h	1.057i
RD S	0.615j	0.427i	1.041j
DCD +S	0.678c	0.488c	1.166c
75% DCD	0.659e	0.469e	1.128e
RD S	0.631h	0.444g	1.075h
DCD +S	0.687b	0.496b	1.183b
100% DCD	0.668d	0.481d	1.149d
RD S	0.642g	0.454f	1.096g
DCD +S	0.694a	0.507a	1.202a
LSD <sub>at 5%</sub>	0.007	0.007	0.010

RD (recommended dose), DCD ( Dicyandiamide), S (sulphure), FW (Fresh weight)

### • N, P, K and S concentration:

Data in Table 2 show that N, P, K and S concentrations in leaves were significantly increased due to supplying the plants with NPK fertilization at different rates. The highest mean values were recorded with adding 100% NPK from recommended dose. As

for the increasing N, P and K concentration with increasing rates of NPK fertilization, it may be owed to the availability of N, P and K elements for plant and improving root growth, hence increasing the absorbing area of root. These results agree with findings of Gairola *et al.*, (2009); Mirdad, (2009); Kawthar *et al.*, (2014); Singh *et al.*, (2014); Hossain *et al.*, (2014) and Nemadozi, (2015) who showed that N, P, K concentrations were increased with increasing rates of NPK fertilization and chlorophyll content.

Data in Table 2 show that the sulphur application + DCD inhibitor had a significant effect on N, P, K and S concentrations in leaves of lettuce. All elements concentrations in lettuce leaves were significantly increased with adding sulphur+DCD during the season. These results may be related to the positive effect of sulphur on the availability of some nutrients, which was reflected on plant uptake and plant growth (Mehana and Farag, 2000). An additional advantage of Urea mix with DCD is improving the uptake of macro and micronutrients. When plant roots take up ammonium ions protons (H<sup>+</sup>) are excreted for charge equilibration in the roots. Thus, the pH of the rhizosphere is decreased, resulting in greater mobility and uptake of some nutrients by the plant. These results are in accordance with the finding of Choudhury and Kennedy (2005), Shaviv (2005), Amberger (2008) and Trenkel (2010).

**Table2. Effect of sulphur application, Dicyandiamide (DCD) and NPK mineral fertilization rates on N, P, K and S% in lettuce leaves.**

Char. Treat.	N%	P%	K%	S%
<b>NPK fertilization</b>				
50% RD	1.97b	0.197c	2.20c	0.251c
75% RD	2.11a	0.217b	2.41b	0.293b
100% RD	2.20a	0.227a	2.51a	0.333a
LSD <sub>at 5%</sub>	0.14	0.05	0.04	0.005
<b>S, DCD and S+DCD fertilization</b>				
DCD	2.02b	0.210b	2.33b	0.224c
S	1.80c	0.187c	2.11c	0.315b
DCD +S	2.44a	0.244a	2.68a	0.338a
LSD <sub>at 5%</sub>	0.13	0.003	0.05	0.005
<b>Interaction</b>				
100% RD	2.01cde	0.208f	2.32e	0.209i
50% RD	DCD	1.77fg	0.183i	2.05h
	S	1.66g	0.173j	1.97i
	DCD +S	2.47a	0.234c	2.59c
75% RD	DCD	2.11cd	0.219e	2.41d
	S	1.82efg	0.189h	2.13g
	DCD +S	2.39ab	0.243b	2.69b
100% RD	DCD	2.20bc	0.227d	2.53c
	S	1.93def	0.200g	2.22f
	DCD +S	2.47a	0.254a	2.77a
LSD <sub>at 5%</sub>	0.20	0.006	0.07	0.008

It is obvious from data in Table 2 that the interaction between S and DCD had significant effects

on all chemical constituents in lettuce under all treatments of NPK rats. On the other hand, the highest values of elements concentration in lettuce are connected with 100% NPK + S + DCD which were 2.47, 0.254, 2.77 and 0.384 for N, P, K and S%, respectively.

**• NO<sub>3</sub>-N, NO<sub>2</sub>-N and nitrate reductases activity:**

Obtained results in Table 3 illustrated the effect of different rates of NPK fertilization, sulphur, DCD and their interactions NO<sub>3</sub>-N, NO<sub>2</sub>-N and nitrate reductases activity during 2016.

**Table3. Effect of sulphur application, Dicyandiamide (DCD) and NPK mineral fertilization rates on NO<sub>3</sub>-N, NO<sub>2</sub>-N and nitrate reductases activity in lettuce leaves.**

Char. Treat.	NO <sub>3</sub> -N mg.l <sup>-1</sup>	NO <sub>2</sub> -N mg.l <sup>-1</sup>	Nitrate reductases activity mg.min .g <sup>-1</sup> FW
<b>NPK fertilization</b>			
50% RD	120.3c	1.04c	2.09a
75% RD	130.6b	1.16b	1.93b
100% RD	138.5a	1.28a	1.75c
LSD <sub>at 5%</sub>	3.34	0.04	0.02
<b>S, DCD and S+DCD fertilization</b>			
DCD	127.9c	1.13b	1.89b
S	160.4b	1.51a	1.39c
DCD +S	101.2a	0.83c	2.49a
LSD <sub>at 5%</sub>	2.66	0.05	0.04
<b>Interaction</b>			
100% RD	181.7a	1.78a	1.09j
50% RD	DCD	116.4g	0.99g
	S	150.5d	1.37d
	DCD +S	93.9j	0.75
75% RD	DCD	128.1f	1.12f
	S	161.8c	1.52c
	DCD +S	102.0i	0.82h
100% RD	DCD	139.2e	1.27e
	S	168.7b	1.64b
	DCD +S	107.6h	0.92g
LSD <sub>at 5%</sub>	4.21	0.08	0.06

It is evident from these results that treating lettuce plants with NPK fertilization significantly increased NO<sub>3</sub>-N and NO<sub>2</sub>-N in lettuce leaves and decreased the nitrate reductases activity. Generally, the most favorable treatment for enhancing NO<sub>3</sub>-N and NO<sub>2</sub>-N mg.l<sup>-1</sup> ,in leaves were the treatment of 100% NPK from RD, while nitrate reductases activity recorded the highest values with 50% from RD. This trend was true during season of the experimentation. Abd-Elfattah, (2012) illustrated that the highest nitrate content due to chemical fertilization maybe attributed to that mineral fertilizer salts are soluble and nitrogen is immediately available for plant uptake soon after fertilizer application.

It is clear from data in Table 3 that the positive effects were happened in NO<sub>3</sub>-N, NO<sub>2</sub>-N mg.l<sup>-1</sup> , and nitrate reductases activity of lettuce with using of DCD,

sulphur, S + DCD. The highest values of NO<sub>3</sub>-N, and NO<sub>2</sub>-N mg.l<sup>-1</sup> were recorded with adding sulphur alone comparing with other treatments while nitrate reductase activity of lettuce was connected with DCD+S. These results could be attributed to the sulfur's role in the activation of the enzyme nitrate reductase, which is necessary for the conversion of nitrate to amino acids in plants. Low nitrate reductase activity depresses soluble protein levels, while raises nitrate concentrations in plant tissue. The obtained results are in harmony with those reported by Cuppett *et al.*, (1999), Al-Redhaiman *et al.*, (2003) Wright *et al.*, (2010) and Sanchez and Silvertooth (2016). Because accumulation of nitrate in plants depended on its concentration in the soil solution; an addition of nitrification inhibitors to ammonical N-fertilizers will decrease the concentration of nitrate in the soil and consequently its contents in the plant. Similar results were reported by Pasad *et al.*, (2001b), Amberger (2006), Shaviv (2005) and Grant (2005).

Data in Table 3 reveal that NO<sub>3</sub>-N, NO<sub>2</sub>-N mg.l<sup>-1</sup> and nitrate reductase activity of lettuce was significantly affected during 2016 with the interaction among the treatments comparing with the control (100% NPK). The lowest NO<sub>3</sub>-N, NO<sub>2</sub>-N mg.l<sup>-1</sup> were obtained by applied S+DCD+ 50% RD from NPK comparing with the control (100% NPK) which recorded the highest mean values. While, nitrate reductase activity of lettuce recorded the highest mean values with S+DCD+ 50% over the control which was 2.63 mg.l<sup>-1</sup>.

• **Available N, P and K mg.l<sup>-1</sup>:**

Data in Table 4 indicate that available N, P and K mg.l<sup>-1</sup> in the soil after harvesting significantly affected with application of NPK fertilization rates during the season of the experiment. Moreover, application of 100% NPK from recommended dose was more effective than the other treatments to increase the average values. The obtained results are in accordance with Titilola (2006); Ayeni and Adetunji (2010) and Onwudiwe *et al.*, (2014).

Data illustrated in Table 4 show the effect of DCD, sulphur, S + DCD on availability of N, P and K mg.l<sup>-1</sup> of lettuce. Such data indicate that the application of sulphur + DCD increased availability of N, P and K mg.l<sup>-1</sup>. The positive effect of this sulphur level may be due to lowering soil pH value, improving soil structure, soil chemical properties and increased the availability of certain plant nutrients such as N, P and K also, several micronutrients i.e. Fe, Zn and Mn. The obtained results are in accordance with those obtained by Al-Redhaiman *et al.*, (2003); Wright *et al.*, (2010) and Sanchez and Silvertooth (2016). Adding of DCD mix with N-fertilization in form of urea improved N, P and K availability because of Protons (H<sup>+</sup>) are excreted for charge equilibration in the roots. Thus, the pH of the rhizosphere is decreased, resulting in greater mobility and elements become available for plant. These results are in accordance with the finding of Amberger (2008) and Trenkel (2010).

It is clear from data in Table 4 that there were significant interactions among treatments under study on availability of N, P and K mg.l<sup>-1</sup> in the soil after

harvesting during season of 2016. In general, plants fertilized by mix of S+DCD produced the highest values of traits under all rates of NPK fertilization comparing with the control (100% NPK). In the same time the highest values of available N, P and K realized with addition of 100% NPK+ S+ DCD over the control which recorded values near to that using DCD + 75% NPK.

**Table4. Effect of sulphur application, Dicyandiamide (DCD) and NPK mineral fertilization rates on N, P and K mg.l<sup>-1</sup> in the soil after harvesting.**

Char. Treat.	N mg.l <sup>-1</sup>	P mg.l <sup>-1</sup>	K mg.l <sup>-1</sup>
NPK fertilization			
50% RD	61.22c	7.60b	253.4b
75% RD	69.09b	8.99c	276.7c
100% RD	72.91a	9.61a	286.8a
LSD <sub>at 5%</sub>	1.39	0.16	2.19
S, DCD and S+DCD fertilization			
DCD	65.90b	8.34c	266.7c
S	58.67c	6.93b	243.6b
DCD +S	79.66a	10.94a	306.7a
LSD <sub>at 5%</sub>	0.90	0.08	3.38
Interaction			
100% RD	66.47f	8.33f	267.3f
DCD	55.30i	6.57i	236.2i
50% RD S	54.90j	5.94j	228.9j
DCD +S	76.47c	10.28c	295.2c
DCD	69.70e	8.96e	277.4e
75% RD S	58.50h	7.09h	245.4h
DCD +S	79.07b	10.93b	307.4b
DCD	72.70d	9.49d	286.5d
100% RD S	62.60g	7.74g	256.4g
DCD +S	83.43a	11.59a	317.4a
LSD <sub>at 5%</sub>	1.49	0.25	3.82

**CONCLUSION**

From the results of this study, it could be concluded that, application of S + DCD mixed with nitrogen fertilization is the recommended treatment for good chemical composition of lettuce. Based on the obtained results, combined application of nitrogen, phosphorous and potassium is the most suitable and recommendable dose for a good chemical composition of lettuce. So adding S + DCD+100% NPK from recommended dose was the best treatment for a good lettuce.

**REFERENCES**

Abd-Elfattah, A. M. (2012). Effect of bio and organic fertilizer and some foliar applications on tomato fruits yield and quality (*Lycopersicon esculentum, Mill*). M. Sc. Thesis, Fac. of Agric. Mans. Univ., Egypt.

Al-Redhaiman, K. N.; M. I. D. Helal and R. R. Shahin (2003). Effect of sulfur blended N-fertilizers on nitrogen use efficiency and quality of lettuce yield. Pakistan J. Biol. Sci., 6 (16): 1408-1412.

- Amberger, A. (2006). Soil Fertility and Plant Nutrition in the Tropics and Subtropics. International Fertilizer Industry Association, Paris, France; International Potash Institute, Horgen, Switzerland.
- Amberger, A. (2008). 100 Jahre Kalkstickstoff – ein multifunktionaler N-Dünger. (German) AlzChem Trostberg GmbH, Chemiepark Trostberg, Germany.
- Asing, J.; S. Saggarr, J. Singh and N. S. Bolan (2008). Assessment of nitrogen losses from urea and an organic manure with and without nitrification inhibitor, dicyandiamide, applied to lettuce under glasshouse conditions. Australian Journal of Soil Research, 46: 535–541.
- Ayeni, L. S. and M. T. Adetunji (2010). Integrated Application of Poultry Manure and Mineral Fertilizer on Soil Chemical Properties, Nutrient Uptake, Yield and growth components of maize. Nature and Sci., 8(1): 60-67.
- Choudhury, A. T. M. A. and I. R. Kennedy (2004). Nitrogen Fertilizer Losses from Rice Soils and Control of Environmental Pollution Problems. Communications in Soil Science and Plant Analysis, 36: 1625 – 1639
- Cui, M.; X. C. Sun, C. X. Hu, H. J. Di, Q. L. Tan and C. S. Zhao (2011). Effective mitigation of nitrate leaching and nitrous oxide emissions in intensive vegetable production systems using a nitrification inhibitor, dicyandiamide. J. Soils Sediment., 11, 722–730.
- Cuppertz, S. L.; M. Mcvey McCluskey, E. T. Paparozzi and A. Parkhurs (1999). Nitrogen and sulfur effects on leaf lettuce quality. J. Food Quality 22: 363-373.
- Di, H. J.; K. C. Cameron and R. R. Sherlock (2007). Comparison of the effectiveness of a nitrification inhibitor, dicyandiamide, in reducing nitrous oxide emissions in four different soils under different climatic and management conditions. Soil Use Manag., 23, 1–9.
- Di, H. J.; K. C. Cameron, J. P. Shen, C. S. Winefield, M. O' Callaghan, S. Bowatte and J. Z. He (2010). Ammonia oxidizing bacteria and archaea grow under contrasting soil nitrogen conditions. FEMS Microbiology Ecology, 72: 386–394.
- Di, H. J.; K. C. Cameron, J. P. Shen, C. S. Winefield, M. O'Callaghan, S. Bowatte and J. Z. He (2011). Methanotroph abundance not affected by applications of animal urine and a nitrification inhibitor, dicyandiamide, in six grazed grassland soils. J. Soils Sediment., 11, 1032–1039.
- El Sayed, H. E. A.; A. H. A. El-Morsy and H. S. Al Othaimen (2015). The Role of Sulphur and Certain Foliar Spray Levels of Micro-nutrients on Garlic (*Allium sativum* L.) Plant. Int. J. Curr. Res. Biosci. Plant Biol. 2015, 2(6): 76-87.
- Gairola, S.; S. Umar and S. Suryapani (2009). Nitrate accumulation, growth and leaf quality of spinach beet (*Beta vulgaris* Linn.) as affected by NPK fertilization with special reference to potassium. Indian J. Sci. and Tech., 2(2): 35-40.
- Gavrilenko, V. F. and T. V. Zigalova (2003). The Laboratory Manual for the Photosynthesis. Academia, Moscow. 256 crp. (in Russian).
- Gomez, K. A. and A. A. Gomez (1984). "Statistical Procedures for Agricultural Research". John Wiley and Sons, Inc., New York. pp:680.
- Grant, C. (2005). Policy aspects related to the use of enhanced-efficiency fertilizers: Viewpoint of the scientific community. IFA International Workshop on Enhanced-Efficiency Fertilizers, Frankfurt. International Fertilizer Industry Association, Paris, France.
- Hageman, R. H. and A. J. Reed (1980). In: Methods in Enzymology. Vo. 69 Part C (Ed Anthony San Pietro) Academic Press New York P 270.
- Haluschak, P. (2006). Laboratory Methods of Soil Analysis. Canada-Manitoba Soil Survey. April
- Hossain, N.; M. Islam, M. Alamgir and M.G. Kibria (2014). Growth Response of Indian Spinach to Biogas Plant Residues. J. Pharmacy and Bio. Sci., 9 (3 & 4):1-6.
- Hosseney, M. H. and M. M. Ahmed (2009). Effect of nitrogen, organic and biofertilization on productivity of lettuce (cv. *Romaine*) in sandy soil under assiat conditions. Ass. Univ. Bull. Environ. Res. 12: 79.
- Jones, D. L.; P. G. Dennis, A. G. Owen and P. A. W. Van Hees (2003). Organic acid behavior in soils - misconceptions and knowledge gaps. Plant and Soils 248: 31 – 41. doi: 10.1023/A:1022304332313.
- Kawthar, A. E.; A. H. M. Rabie and H. H. Manaf (2014). Influence of Nitrogen Forms on Nitrate and Nitrite Accumulation in the Edible Parts of Spinach (*Spinacia oleracea*, L.) Plant with Maintenance for Yield Production. J. Horti. Sci. and Ornamental Plants. 6 (3): 126-132.
- MALR (Ministry of Agriculture and Land Reclamation) (2012). Economic Affairs Sector. The Indicators Agriculture Statistics 2:219.
- Masarirambi, M. T.; P. Dlamini, P. K. Wahome and T. O. Oseni (2012). Effects of Chicken Manure on Growth, Yield and Quality of Lettuce (*Lactuca sativa* L.) 'Taina' Under a Lath House in a Semi-Arid Sub-Tropical Environment. American-Eurasian J. Agric. & Environ. Sci., 12:399.
- Mehana, T. A. and F. M. Farag (2000). Influence of phosphate-dissolving micro organisms and elemental sulphur on phosphorus and micronutrient availability in a calcareous soil treated with rock phosphate. J. Agric. Sci. Mansoura Univ. 25(5), 2983-2993.
- Mertens, D., (2005a). AOAC official method 922.02. Plants preparation of laboratory sample. Official methods of analysis, 18<sup>th</sup> edn. North Frederick Avenue, Gaitherburg, Maryland, pp.1-2
- Mertens, D., (2005b). AOAC Official method 975.03. Metal in plants and pet foods. Official methods of analysis, 18<sup>th</sup> edn. North Frederick Avenue, Gaitherburg, Maryland, pp. 3-4

- Midan, S. A. and M. E. Sorial (2011). Some Antioxidants Application in Relation to Lettuce Growth, Chemical Constituents and Yield. Australian J. of Basic and Appl. Sci., 5:127.
- Mirdad, Z. N. (2009). Spinach (*Spinacia oleracea*, L.) Growth and yield responses to irrigation dates, mineral nitrogen - sources and levels - application. J. Agric. & Env. Sci. Alex. Univ., Egypt. 8 (1): 43-69.
- Morris, R. G.; S. L. Tisdal and J. Platou (1984). The importance of sulphur in crop quality. J. Fert. Issues 1, 139-145.
- Nabih, M. A. (2002). Studies on potatoes (*solanum tuberosum* L.). M. Sc. Thesis, Fac. of Agric. Mans. Univ., Egypt.
- Nemadodzi, L. E. (2015). Growth and development of baby spinach (*Spinacia oleracea* L.) with reference to mineral nutrition. M.Sc. Thesis, Fac. Sci., University of South Africa.
- O'Callaghan, M.; E. M. Gerard, P. E. Carter, and R. Lardner (2010). Effect of the nitrification inhibitor dicyandiamide (DCD) on microbial communities in a pasture soil amended with bovine urine. Soil Biol. Biochem., 42, 1425-1436.
- Okoli, P. S. O and I. A. Nweke (2015). Effect of poultry manure and mineral fertilizer on the growth performance and quality of cucumber fruits. J. of Experimental Biology and Agric. Sci., 3(4):362-367.
- Onwudiwe, N.; O. U. Benedict; P. E. Ogbonna and E. E. Ejiofor (2014). Municipal solid waste and NPK fertilizer effects on soil physical properties and maize performance in Nsukka, Southeast Nigeria. African J. Biotech., 13(1): 68-75.
- Reeuwijk, L. P. (2002). Procedures For Soil Analysis. Inter. Soil Ref. and Info. Center. Food and Agric. Organization of the United Nations
- Sahlemedhin, S. and B. Taye (2000). Procedures for Soil an Plant Analysis. Technical Paper No. 74.
- Saif El-Din, U. Z. (2000). Studies on bio and chemical fertilization on sweet potato (*Ipomoea batatas* L.). M. Sc. Thesis, Fac. Agric. Mans. Univ., Egypt.
- Sanchez, C. A. and J. C. Silvertooth (2016). Evaluation of Soil Amendments for Lettuce Production in the Desert. College of Agric., Univ. Arizona (Tucson, AZ): 51-62.
- Shaviv, A. (2005). Controlled Release Fertilizers. IFA International Workshop on Enhanced-Efficiency Fertilizers, Frankfurt. International Fertilizer Industry Association Paris, France.
- Singh, J. P. (1988). A rapid method for fertermination of nitrate in soil and plant extracts. Plant and soil. 110: 137-139.
- Singh, S.; M. Agrawal and S.B. Agrawal (2014). Differences in responses of summer and winter spinach to elevated UV-B at varying soil NPK levels. Environ Sci Pollut Res (2014) 21:6099-6107.
- Titilola, A. O. (2006). Effects of Fertilizer Treatments on Soil Chemical Properties and Crop Yields in a Cassava-based Cropping System. J. Applied Sci. Res., 2(12): 1112-1116.
- Trenkel, M. E. (2010). "Slow- and Controlled-Release and Stabilized Fertilizers: An Option for Enhancing Nutrient Use Efficiency in Agriculture". Paris, France.
- Wright, A. L.; Y. Luo; N. Havranek; D. S. David; W. R. Ronald and Y. Rongzhong (2010). Elemental Sulfur Use for Increasing Phosphorus Availability to Lettuce in Everglades Agricultural Area Soil. A Horticultural Sciences Department Extension Publication on Vegetable and Fruit Crops.

## دور اضافة الكبريت والمثبط مقارنة بالتسميد المعدني على التركيب الكيماوي لنبات الخس كريم فكري فودة قسم الأراضي - كلية الزراعة - جامعة المنصورة - مصر .

اجريت تجربه حقلية لدراسة تأثير اضافة الكبريت مع المثبط مقارنة باضافة التسميد المعدني على التركيب الكيماوي لنبات الخس خلال تجربه عاملية تشتمل على 9 معاملات وهي عبارة عن 3 مستويات من التسميد المعدني (50، 75، 100 % من الموصي به) و 3 معاملات من (المثبط، كبريت، مثبط + كبريت) خلال 3 مكررات وبالتالي يصبح المجموع الكلي 27 وحده تجريبية مقارنة بمعامله كنترول (100 % موصي به من التسميد المعدني فقط). اوضحت النتائج ان محتوى الكلوروفيل ، النسبة المئوية للنيتروجين، الفوسفور ، البوتاسيوم ، الكبريت في الخس و النترات والنيتريت بالاضافة الى صلاحية النيتروجين ، الفوسفور والبوتاسيوم في التربة بعد الزراعة تآثرت معنويا باضافة مستويات التسميد المعدني وسجلت اعلى القيم عند استخدام 100 % من الموصي به بينما سجلت اعلى القيم نشاط انزيم النيتريت انخفضت مع استخدام التسميد المعدني . و كانت اعلى القيم السابقة عند استخدام خليط من المثبط مع الكبريت مقارنة بباقي المعاملات ماعدا النترات والنيتريت سجلت اعلى القيم عند استخدام الكبريت فقط. في نفس الاتجاه الاضافة المشتركة بين المعاملات تحت الدراسة على نفس الصفات اوضحت ان اعلى القيم سجلت عند استخدام خليط الكبريت و المثبط مع 100 % تسميد معدني من الموصي به بينما نشاط انزيم النيتريت سجلت عند استخدام نفس الخليط عند 50 % من التسميد المعدني مقارنة بمعامله الكنترول.