

## **RESPONSE OF EGYPTIAN HYBRID1 RICE TO ORGANIC AND INORGANIC SOURCES OF NITROGEN FERTILIZER**

**Abdel-Fattah, G. A. ; W. M. El Khoby, A. M. El-Ekhtyar and B.B. Mikhael**

**Rice Res. and Training Center, Field Crops Res. Inst., Agric. Res. Center, 33717 Sakha-Kafr El-Sheikh, Egypt**

### **ABSTRACT**

Low levels of available nitrogen in soils may limit rice growth. An investigation was undertaken at the farm of Rice Research and Training Center, Sakha, Kafr El-Shiekh during 2010 and 2011 rice growing seasons the objectives of this investigation was aimed to evaluate the effect of different nitrogen sources combination on growth, yield and its components of Egyptian hybrid rice 1 (EHR1). All growth traits i.e. plant height, number of tillers hill<sup>-1</sup>, chlorophyll content as well as dry matter accumulation differed significantly by N sources combination. Application of 46 kg N fed<sup>-1</sup> plus 7 tons of FYM fed<sup>-1</sup> recorded highest values of these traits at all different growth stages, while the untreated control treatment gave the lowest ones. Applications of 46 kg N fed<sup>-1</sup> plus 7 tons FYM fed<sup>-1</sup> produced tallest panicles and highest panicle grain weight, number of panicles hill<sup>-1</sup>. The untreated (control) treatment gave the minimum values of all traits mentioned above.

**Keywords:** Hybrid Rice, FYM, Composted Rice Straw, Mineral nitrogen

### **INTRODUCTION**

Rice (*Oryza sativa* L.) is one of the most important cereal crops of the world, grown in a wide range of climatic zones, to nourish the mankind. Introduction of hybrid rice is an important step towards augmentation by about 15-20% more than the promising high-yielding commercial varieties. Earlier studies revealed that judicious and proper use of fertilizers can markedly increase the yield and improve the quality of rice (Chaturvedi 2005). Increasing rice production can be achieved also through improving management systems of crop culture, especially the nutrient management of the crop as well as the proper utilization of the different sources of nutrients, i.e. the natural nutrient reserve of soil, chemical, organic and bio-fertilizer. The nutrient management aims to reduce agrochemical use and enhance soil fertility through using different sources of nitrogen fertilizer.

None of such sources is complete or sufficient to sustain soil fertility and crop productivity; hence the growers are obliged to use more levels of inorganic chemical fertilizers. Generally, urea is the most convenient N source for rice. The efficiency of the urea-N in rice culture is low, around 30-40%, in some cases even lower (Choudhury and Kennedy 2005).

The application of the organic fertilizers such as the farmyard manure, compost and green manure, increase the organic matter content which serves neural advantages like conservation and slow release of

nutrients (Chaturvedi 2005). These advantages lead to increasing the fertility and productivity of the soil (Choudhury and Kennedy 2005). Nitrogen fertilizer losses through different mechanisms can be minimized by reducing the amount of applied N fertilizer with an efficient use of N by the rice plant. Plant growth promoting microorganisms can reduce the use of urea-N by growth promotion through the production of auxines, cytokinins, gibberellins, and ethylene (Dobbelaere *et al.* 2003). Wide range of biological nitrogen fixing agents (BNF agents) i.e. Cyanobacteria and Azospirillum spp. (bacteria are indigenous and can be growth in the flooded rice fields where sufficient water and an aerobic condition are favorable to their growth. The BNF technology are considered important for long term maintenance of soil fertility, economically justifiable and environmentally safe improving soil properties, reducing pests and diseases and reducing environmental pollutions (Choudhury and Kennedy 2005). The replenishment can be attained using a combination of different sources of N fertilizer. Roger (1995) recorded an average increase in yield of 19.8 % due to bacterial inoculation of rice. Similar trend was obtained for harvest index value. Also, N fertilizer efficiency in control plots was 18.7 Kg grain/ kg N applied while for inoculated plots was 19.1 kg grain/ kg N applied. El-Hawary and Hammodua (1986) found that application of certain partially decomposed organic manure, gave significant increase in rice grain yield. Also, they reported that application of these amendments with *Azotobacter* spp, gave higher yield compared to their application without bacteria.

Annually, the amount of agricultural waste produced in Egypt is enormous especially rice straw. The effect of using agricultural waste composts to fertilize agricultural land has been positive from the perspective of a recycling economy and because of their valuable characteristics and ingredients (Eneji, *et al.* 2001). Baziramakenga and Simard (2001) reported that organic materials contain many essential elements at low concentrations, which are slowly released upon decomposition. Manure is used mainly as a source of nitrogen (N) and phosphorus (P) (Materechera and Salagae 2002). The soil pH, organic carbon, and available P and potassium (K) concentrations of soils increase with the application of compost. Application of manure enhances rice nutrient accumulation and dry matter yield (Eneji, *et al.* 2001).

The rice stem borer infection in rice plants was found negatively correlated with high soil and plant silica (Djamin and Pathak 1967, Rani *et al* 2006). The positive effect of applying Azospirillum in raising phenolic levels, that reduce insect infestation in rice, was reported by (Mohan *et al* 1988).

The objective of the present study is to evaluate Egyptian hybrid rice 1 (EHR1) response to different nitrogen source combinations. Also, the level of rice stem borer damage as affected by fertilizer combination was investigated.

## MATERIALS AND METHODS

The present investigation was conducted at the farm of Rice Research and Training Center Sakha, Kafr El- Shiekh, Egypt during 2010 and 2011 rice summer seasons the objective of this investigation was aimed to study the effect of chemical N, organic and bio-N fertilizer and their combinations on the productivity of Egyptian hybrid rice 1. The inorganic fertilizer as urea (46.5% N), farmyard manure as cattle manure and bacterial inoculation as *Azospirillum* spp as commercial name Cerialen were used. The nitrogen was applied as ten treatments (T<sub>0</sub> control, (T<sub>2</sub>) 69 kg N fed<sup>-1</sup> as urea,(T<sub>3</sub>) 7 t FYM fed<sup>-1</sup>., (T<sub>4</sub>) 2 tons composted rice straw fed<sup>-1</sup>. (T<sub>5</sub>) Bacterial inoculation (BI) 400 gm Cerialen fed<sup>-1</sup>., (T<sub>6</sub>) 46 kg N fed<sup>-1</sup> + 7 t FYM fed<sup>-1</sup>, (T<sub>7</sub>) 46 kg N fed<sup>-1</sup> + 2 tons composted rice straw fed<sup>-1</sup>, (T<sub>8</sub>) 46 kg N fed<sup>-1</sup> + BI, (T<sub>9</sub>) 7 t FYM fed<sup>-1</sup> + BI, (T<sub>10</sub>) 2 tons composted rice straw fed<sup>-1</sup> + BI. The mechanical and chemical soil analysis is presented in Table 1, and the chemical analysis of compost and farm yard manure which used in this trial are shown in Table (2).

**Table 1. Some physical and chemical properties of soil of the experimental location.**

Soil properties	2010	2011
Mechanical analysis		
Ca <sup>++</sup>	9.5	10.0
Mg <sup>++</sup>	3.94	3.98
K <sup>+</sup>	1.76	1.80
Na <sup>+</sup>	14.8	15.2
Soluble anions, megl <sup>-1</sup>		
CO <sub>3</sub> <sup>=</sup>	0.00	0.00
HCO <sub>3</sub> <sup>-</sup>	6.00	6.75
Cl <sup>-</sup>	8.30	8.44
SO <sub>4</sub> <sup>=</sup>	15.7	15.79
Available micronutrients, ppm		
Fe <sup>++</sup>	6.10	5.80
Zn <sup>++</sup>	1.10	1.05
Mn <sup>++</sup>	3.50	3.01

The complete randomized block design with four replication was used. Each sub-plot measured 3 m in width and 5 m in length. All the recommended cultural practices were followed for seedbed preparation as well as for permanent field. The decomposed farm yard manure and compost fertilizer were applied in dry soil, then incorporated well during tillage, while two third of inorganic N fertilizer was applied two third before flooding and one third at panicle initiation stage. The inoculation was performed according to Omer *et al.* 1989.

**Table 2. Chemical components of the organic matter**

Nutrient	FYM		Composted rice straw	
	2010	2011	2010	2011
<b>C%</b>	34.4	34.8	28.5	29.13
<b>N%</b>	1.30	1.42	1.83	1.92
<b>C/N Ratio</b>	26.70	27.16	15.57	16.26
<b>P%</b>	0.38	0.43	2.30	2.55
<b>K%</b>	0.39	0.46	2.43	2.64
<b>Fe ppm</b>	600	633	510	560
<b>Mn ppm</b>	300	340	490	489
<b>Zn ppm</b>	69	58	54	61

For the bacterial inoculation treatment in the nursery, the pre-germinated seeds were mixed with the bacterial (*Azospirillum*) suspension overnight just before sowing as powdered inoculums (water adhesive like sugar or gum). Thus, each seed received about  $10^8$  bacterial cells. Then, the seeds were broadcasted by hand on May 5<sup>th</sup> and May 9<sup>th</sup> in 2010 and 2011 seasons, respectively. In permanent field, at transplanting the seedlings were inoculated again by soaking their roots over night in the corresponding bacterial suspension to complete the required rate of bacterial fertilizers. To determine the growth characters at maximum tillering and panicle initiation stages, three hills per plot were harvested and the number of tillers hill<sup>-1</sup> was recorded, chlorophyll content was measured using chlorophyll-Meter, and plant height was measured then it was dried and weighed to determine the total dry weight. At harvest, average number of tillers, plant height and panicle length, panicle characters, grain and straw yield were estimated in each plot, and grain yield was calibrated to 14% moisture basis.

The collected data for each trait were subjected to the analysis of variance according to the procedure outlined by Gomez and Gomez (1984). Differences among treatment means were compared using the Revised L.S.D at 5% levels of significance adopted by Waller and Duncan (1969). The homogeneity test was conducted of error mean squares and, accordingly, the combined analysis was carried out between years because such errors were homogeneous.

## RESULTS AND DISCUSSION

### A. plant growth parameters.

The fertilizer treatments had significant effect on the plant growth at different growth stages (Tables 3, 4 and 5). Plant height increased gradually with increasing plant age. All nitrogen treatments caused significant increases in plant height over control. The application of 46 kg N as urea + 7 t FYM fed<sup>-1</sup> recorded highest unfavorable value of plant height at different growth stages, while the application of 400 g Cerialen alone gave the shortest plants. The increase in plant height in response to application of N fertilizers is probably due to enhanced availability of nitrogen which enhanced leaf area resulting in higher photo assimilates

and consequently more dry matter accumulation. These results are similar to the findings of Mandal *et al.* (1992) and (Chaturvedi 2005).

**Table 3. Rice growth characters of hybrid 1 at maximum tillering stage as affected by different combination of nitrogen sources.**

Treatment	Plant height cm		Number of tillers hill <sup>-1</sup>		Chlorophyll content (SPAD)		Dry matter accumulation g hill <sup>-1</sup>	
	2010	2011	2010	2011	2010	2011	2010	2011
Control	56.0	55.49	20.50	19.77	30.41	29.84	22.17	21.94
69 kg N fed <sup>-1</sup> Urea	82.50	81.81	32.75	31.21	41.54	40.10	44.12	42.96
7 t FYM fed <sup>-1</sup>	73.0	71.31	28.65	27.12	41.04	39.06	42.25	40.38
2 t composted rice straw fed <sup>-1</sup>	61.50	60.49	25.44	24.36	38.11	37.58	35.95	35.24
400 gm Cerialen	62.90	63.20	24.66	23.51	39.22	38.19	31.17	29.48
46 kg N fed <sup>-1</sup> Urea + 7 t FYM fed <sup>-1</sup>	87.0	86.41	34.18	33.11	42.17	40.61	59.22	57.94
46 kg N fed <sup>-1</sup> Urea + 2 t composted rice straw fed <sup>-1</sup>	83.22	82.12	28.11	27.00	38.89	38.63	56.96	55.67
46 kg N fed <sup>-1</sup> Urea + 400 gm Cerialen	81.32	80.22	29.78	29.14	40.15	39.60	58.32	57.96
7 t FYM fed <sup>-1</sup> + 400 gm Cerialen	73.50	73.37	32.19	31.86	41.16	40.83	43.26	42.78
2 t composted rice straw fed <sup>-1</sup> + 400 gm Cerialen	56.12	55.12	20.13	19.26	37.10	36.89	37.43	36.35
L.S.D. 0.05	3.21	3.18	1.53	1.53	1.19	1.18	2.64	2.52

**Table 4: Rice growth characters of hybrid 1 at complete heading stage as affected by different combination of nitrogen sources.**

Treatment	Plant height cm		Number of tillers hill <sup>-1</sup>		Chlorophyll content SPAD		Dry matter accumulation g/ hill <sup>-1</sup>	
	2010	2011	2010	2011	2010	2011	2010	2011
Control	86.19	84.29	19.10	17.91	35.79	35.55	64.50	62.25
69 kg N fed <sup>-1</sup> Urea	98.95	97.93	29.17	28.51	45.55	46.01	165.29	161.42
7 t FYM fed <sup>-1</sup>	99.28	98.00	23.98	23.85	45.50	44.26	159.10	152.03
2 t composted rice straw fed <sup>-1</sup>	93.72	92.54	20.50	19.43	42.00	41.35	117.55	114.36
400 gm Cerialen	92.16	89.14	21.13	19.71	45.61	44.51	122.18	120.33
46 kg N fed <sup>-1</sup> Urea + 7 t FYM fed <sup>-1</sup>	107.11	106.75	30.44	29.21	48.89	49.06	199.76	200.41
46 kg N fed <sup>-1</sup> Urea + 2t composted rice straw fed <sup>-1</sup>	99.32	97.01	25.44	25.29	45.82	44.62	165.10	162.31
46 kg N fed <sup>-1</sup> Urea + 400 gm Cerialen	106.25	105.63	28.00	27.43	46.87	46.60	171.00	169.90
7 t FYM fed <sup>-1</sup> + 400 gm Cerialen	98.10	96.40	28.66	27.29	46.32	45.28	180.12	177.36
2 t composted rice straw fed <sup>-1</sup> + 400 gm Cerialen	92.44	90.66	17.92	18.30	45.69	44.65	122.15	118.35
L.S.D. 0.05	2.89	2.82	1.69	1.61	1.75	1.66	3.84	3.66

Number of tillers hill<sup>-1</sup> at different growth stages as affected by different combinations of N sources is presented in Table 4. With increasing the plant age, number of tillers were decreased. All N treatments affected significantly number of tillers at all studied growth stages. Application of 46 kg N ha<sup>-1</sup> as urea plus 7 t FYM fed<sup>-1</sup> produced the highest number of tillers hill<sup>-1</sup> followed by the application of 69 kg N fed<sup>-1</sup> urea alone, while was recorded the lowest in nonfertilized plots. More number of tillers m<sup>-2</sup> might be due to the more availability of nitrogen that plays a vital role in cell division. These results are in accordance to the findings of Chaturvedi (2005).

**Table 5. Rice growth characters of hybrid 1 at harvest stage as affected by different combination of nitrogen sources.**

Treatments	Plant height		Number of tillers		Dry matter accumulation	
	cm		hill <sup>-1</sup>		g/ hill <sup>-1</sup>	
	2010	2011	2010	2011	2010	2011
Control	87.11	86.71	16.44	14.06	86.71	84.34
69 kg N fed <sup>-1</sup> Urea	104.50	104.74	28.16	26.38	185.40	180.82
7 t FYM fed <sup>-1</sup>	97.23	97.05	24.00	22.51	177.19	174.04
2tcomposted rice straw fed <sup>-1</sup>	93.13	92.54	19.11	17.09	131.14	126.45
400 gm Cerialen	90.21	89.99	18.55	17.14	142.60	140.5
46 kg N fed <sup>-1</sup> Urea + 7 t FYM fed <sup>-1</sup>	105.10	104.52	26.50	25.18	231.13	228.04
46 kg N fed <sup>-1</sup> Urea + 2 t composted rice straw fed <sup>-1</sup>	97.21	96.82	24.56	23.99	198.22	194.81
46 kg N fed <sup>-1</sup> Urea + 400 gm Cerialen	103.60	103.19	26.17	25.43	197.00	194.82
7 t FYM fed <sup>-1</sup> + 400 gm Cerialen	98.65	98.09	27.91	26.84	203.11	199.65
2 t composted rice straw fed <sup>-1</sup> + 400 gm Cerialen	93.00	92.11	17.83	16.76	150.13	145.34
L.S.D. 0.05	2.78	2.77	1.41	1.31	4.89	4.32

For chlorophyll content, results in Tables 3 and 4 showed that all N treatments recorded higher significant chlorophyll values over control. At maximum tillering stage, the application of 7 t FYM fed<sup>-1</sup> + 400 g Cerialen produced highest chlorophyll content and without any significant differences with application of either 46 kg N fed<sup>-1</sup> Urea + 7 t FYM fed<sup>-1</sup> or 69 kg N fed<sup>-1</sup> Urea. At complete heading stage, application of 46 kg N fed<sup>-1</sup> Urea + 7 t FYM fed<sup>-1</sup> produced the highest values of chlorophyll content. Dry matter accumulation increased significantly with N fertilizer application in rice at all the growth stages of the crop. The results presented in Table ( 6) revealed a significant increase due to nitrogen fertilizer; all N treatments produced more dry matter significantly than control. Also, dry matter increased with increasing plant age. The highest dry matter accumulation was obtained when plants was received 46 kg N as urea plus 7 t FYM fed<sup>-1</sup> at the three growth stages.

**White head percentage**

Results presented in Table (6) show that rice plots fertilized with mineral urea-nitrogen had the highest levels of white heads. 12.4,10.9 % (in 69kg) ,11.2,10.5% (in 46kg urea+7 tFYM) 12.7,11.4% (in 46kg urea+2t compost) in 2010 and 2011 rice seasons respectively . When urea(46 kg N/fed) was combined with FYM (7t/fed), the rice plants suffered 11.2 ,10.5% white heads , but decreased to 8.0, 8.9%white heads when the same level of urea (46 kg N/fed) was combined with 400g Cerialen (*Azospirillum* sp), in the first and second seasons , respectively. Applications of sole FYM (7 t/fed), composted rice straw (2 t/fed) or Cerialen (400 g/fed) induced low stem borer infestations; 5.2, 5.3, 5.6, 5.2 and 5.9, 5.4% white head in the first and second seasons, respectively.

The low rice stem borer infestation in plots treated with composted rice or with farmyard manure could be attributed to the role of silica (high in both treatments) in controlling the rice stem borer. In such concern, Djamin and pathak (1967) found that the incisor region of the mandibles of stem borer larvae fed on rice plants with high Si content were more damaged. Chandraman *et al* (2010) recorded a significant negative correlation between insect incidence (from which is the rice stem borer) and rice plant content of silica. In case of the rice stem borer, the correlation value was calculated as -0.930. Rani *et al* (2006) indicated that application of *Azospirillum* may activate the phenyl ammonia lyase enzyme implicated in biosynthesis of phenolics, resulting in increased plant phenolics that preven insect damage. Similar results were reported by Mohan *et al* (1988).

**Table6: White head percentage of Egyptian hybrid rice 1 as affected by different combinations of nitrogen sources.**

Treatment	White head%	
	2010	2011
Control	10.14	10
69 kg N fed <sup>-1</sup> Urea	12.4	10.9
7 t FYM fed <sup>-1</sup>	5.2	5.3
2 t composted rice straw fed <sup>-1</sup>	5.6	5.2
400 gm Cerialen	5.9	6.4
46 kg N fed <sup>-1</sup> Urea + 7 t FYM fed <sup>-1</sup>	11.2	10.5
46 kg N fed <sup>-1</sup> Urea + 2 t composted rice straw fed <sup>-1</sup>	12.7	11.4
46 kg N fed <sup>-1</sup> Urea + 400 gm Cerialen	8.0	8.9
7 t FYM fed <sup>-1</sup> + 400 gm Cerialen	8.7	8.2
2 t composted rice straw fed <sup>-1</sup> + 400 gm Cerialen	6.2	6.9
<i>L.S.D. 0.05</i>	2.9	2.8

**Yield attributes**

Yield attributes i.e. panicle length cm, no. of panicles hill<sup>-1</sup>, panicle weight, 1000-grain weight, number of Filled grains panicle<sup>-1</sup>, as well as grain yield t fed<sup>-1</sup> are presented in Tables (7&8). Application of 46 kg N fed<sup>-1</sup> as urea plus 7 t FYM fed<sup>-1</sup> produced the tallest panicle and with par

to the applications of 69 kg N fed<sup>-1</sup> alone or 7 t FYM fed<sup>-1</sup> and those treatments surpassed other treatments significantly, while the shortest panicle was obtained with control.

Concerning number of panicles hill<sup>-1</sup>, application of 7 t FYM fed<sup>-1</sup> + 400 g Cerialen produced highest number of panicles without significant with the values obtained from applying either 69 kg N fed<sup>-1</sup> or 46 kg N fed<sup>-1</sup> plus both 7 t FYM fed<sup>-1</sup> and 400 g Cerialen. All N treatments gave the significant values compared with the control. The lowest values were obtained when the plants was fertilized with compost and Bacteria combination.

**Table 7. Panicle length, No. of panicle hill-1 and panicle weight of Egyptian hybrid rice 1 as affected by different combinations of nitrogen sources.**

Treatment	Panicle length (cm)		No. of panicle/ hill <sup>-1</sup>		Panicle weight (g)	
	2010	2011	2010	2011	2010	2011
Control	19.98	20.57	14.12	13.02	2.80	2.82
69 kg N fed <sup>-1</sup> Urea	23.19	24.12	25.44	25.07	4.37	4.41
7 t FYM fed <sup>-1</sup>	24.50	24.02	22.46	22.31	4.32	4.28
2 t composted rice straw fed <sup>-1</sup>	21.10	21.14	16.54	16.71	3.41	3.37
400 gm Cerialen	21.88	21.95	17.10	16.67	3.55	3.65
46 kg N fed <sup>-1</sup> Urea + 7 t FYM fed <sup>-1</sup>	24.50	24.48	24.85	24.62	4.65	4.76
46 kg N fed <sup>-1</sup> Urea + 2 t composted rice straw fed <sup>-1</sup>	23.38	23.46	23.95	23.74	4.33	4.25
46 kg N fed <sup>-1</sup> Urea + 400 gm Cerialen	22.11	22.08	25.12	24.21	3.67	3.78
7 t FYM fed <sup>-1</sup> + 400 gm Cerialen	23.66	23.74	25.66	25.21	4.20	4.23
2 t composted rice straw fed <sup>-1</sup> + 400 gm Cerialen	22.29	22.25	17.10	16.04	3.68	3.89
L.S.D. 0.05	0.50	0.51	1.20	1.123	0.14	0.16

For panicle weight, all N treatments gave significant values compared with control which recorded the lowest value of panicle weight. Increase in panicle weight due to nitrogen treatments might be primarily due to increase in chlorophyll concentration which led to higher photosynthetic rate and ultimately plenty of photosynthesis available during grain development.

The average of both seasons indicated that the number of filled grains panicle<sup>-1</sup> was significantly affected by the different nitrogen treatments. The application of 46 kg N fed<sup>-1</sup> Urea + 7 t FYM fed<sup>-1</sup> produced significantly higher number of filled grains panicle<sup>-1</sup> than the other N treatments and the control (Table 7). The more number of filled grains per panicle was probably due to better nitrogen status of plant during panicle growth period.

For 1000 grain weight, significant differences were detected among all N treatments. Application of 69 kg N Urea produced heaviest 1000-grain weight, while application of 46 kg N as Urea mixed with 2 t



composted rice straw fed<sup>-1</sup> gave the lowest value of this trait.

Grain yield tons fed<sup>-1</sup> as affected by N sources combination is presented in Table 8.

There were significant increases in grain yield with applying different N sources. As a logic result, applying 46 kg N fed<sup>-1</sup> as Urea plus 7 t FYM fed<sup>-1</sup> produced significantly maximum grain yield/fed followed by 46 kg N fed<sup>-1</sup> as Urea + 400 gm Cerialen, then 69 kg N fed<sup>-1</sup> as urea followed by applying 7 t FYM fed<sup>-1</sup> plus 400 gm Cerialen at the same level of significant. It signifies that hybrid rice variety 'Egyptian hybrid rice 1 is adequately stable concerning its genetic potential for yield of rice. There is a very close relation between the yield and its components, especially with number of filled grains per panicle. The improved growth attributes, viz plant height, chlorophyll content and dry-matter production might be responsible for improved yield attributes. It was found that application of nitrogen improves various crop parameters like panicle length, more productive tillers, number of filled grains per panicle and 1000-grain weight thus resulting in higher yields (Chaturvedi 2005).

**Table 8: No of filled grain panicle<sup>-1</sup>, 1000- grain weight g and grain yield t fed.<sup>-1</sup> of Egyptian hybrid rice 1 as affected by different combinations of nitrogen sources.**

Treatment	No of Filled grains/ panicle <sup>-1</sup>		1000 grain weight (g )		Grain yield t fed. <sup>-1</sup>	
	2010	2011	2010	2011	2010	2011
Control	115.00	115.16	21.57	21.50	2.81	2.72
69 kg N fed <sup>-1</sup> Urea	181.77	184.91	22.78	22.92	4.65	4.60
7 t FYM fed <sup>-1</sup>	190.12	190.04	20.55	20.49	4.41	4.03
2 t composted rice straw fed <sup>-1</sup>	160.13	159.13	21.89	21.77	3.31	3.23
400 gm Cerialen	160.56	161.86	21.10	20.70	3.28	3.21
46 kg N fed <sup>-1</sup> Urea + 7 t FYM fed <sup>-1</sup>	207.29	205.09	21.79	21.70	4.98	4.93
46 kg N fed <sup>-1</sup> Urea + 2 t composted rice straw fed <sup>-1</sup>	192.43	188.31	20.50	20.02	4.51	4.32
46 kg N fed <sup>-1</sup> Urea + 400 gm Cerialen	167.18	173.73	22.44	22.15	4.82	4.75
7 t FYM fed <sup>-1</sup> + 400 gm Cerialen	189.17	183.89	22.42	22.31	4.63	4.55
2 t composted rice straw fed <sup>-1</sup> + 400 gm Cerialen	166.11	164.00	22.23	21.98	3.25	3.16
<i>L.S.D. 0.05</i>	3.38	3.40	1.59	1.56	0.49	0.44

## CONCLUSION

It could be concluded that utilization of 46 kg N fed<sup>-1</sup> as Urea in combination with 7 t FYM fed<sup>-1</sup> is more favorable, most efficient and best economically way for rice production under the present experimental conditions for increase rice grain yield and maintain the soil fertility.

However to produce the organic rice without reduction in grain yield, utilization of 7 tons FYM fed<sup>-1</sup> plus 400 gm Cerialen have to be applied.

## REFERENCES

- Baziramakenga, R. and R.R. Simard,. (2001). Effect of deinking paper sludge compost on nutrient uptake and yields of snap bean and potatoes grown in rotation. *Compost Science and Utilization*, 9: 115-126.
- Chandramani P, R, Rajendran, C, Muthiah and C, Chinniah (2010). Organic source induced silica on leaf folder, stem borer and gall midge population and rice yield. *Journal of Biopesticides*, 3 (2): 423-427.
- Chaturvedi, Indira. (2005). Effect of nitrogen fertilizers on growth, yield and quality of hybrid rice (*oryza sativa*). *Journal of Central European Agriculture*. 6(4):611-618.
- Choudhury, A. T. M. A. and I. R. Kennedy (2005). Nitrogen Fertilizer Losses from Rice Soils and Control of Environmental Pollution Problems. *Communications in Soil Science and Plant Analysis*, 36: 1625-1639.
- Dobbelaere, S., J. Vanderleyden, and Y. Okon (2003). Plant growth promoting effects of diazotrophs in the rhizosphere. *Critical Rev. Plant Sci.*, 22: 107-149.
- Djain, A. and M. D. Pathak (1967). Role of silica in resistance to Asiatic rice orer, *Chilo suppressalis*(Walker) in rice varieties. *Journal of Economic Entomology*, 60 : 347 – 351.
- El-Hawarry, F.I;F and M. Hamouda(1986). Effect of certain organic amendments on the efficiency of inoculation with *Azotobacter ssp*. *J.Agric.Sci. Mansura Univ.* 13(2) 629-634.
- Eneji, A.E., S.Yamamoto, and T. Honna(2001). Rice growth and nutrient uptake as affected by livestock manure in four Japanese soils. *Journal of Plant Nutrition*, 24: 333-343.
- Gomez, K.A. and A.A. Gomez (1984). *Statistical procedure for agricultural research*. 2<sup>th</sup> ed. John Wiley Sons. New yourk. USA.
- Mandal, N.N., P.P. Chaudhry, and D. Sinha(1992). Nitrogen, phosphorus and potash uptake of wheat (var. Sonalika). *Env. and Eco.* 10: 297 (Field Crop Abst. 46 (1): 30).
- Materechera, S.A. and A.M. Salagae(2002). Use of partially decomposed cattle and chicken manure amended with wood ash in two South African arable soils with contrasting texture: Effect on nutrient uptake, early growth, and dry matter yield of maize. *Communications in Soil Science and Plant Analysis*, 33: 179-201.
- Mohan S, D, Purushothaman, S, Jayaraj and A.V.Rangarajan ( 1988). PALase activity in the roots of *Sorghum bicolor* (L.) inoculated with *Azospirillum*. *Curr. Sci.* 57:492-493.
- Omer ,N ;T.Heulin ; P.Weinhord ; M.N.Alaa El Deen and J.Balandreau( 1989). Field inoculation of rice with vitro selected plant growth promoting rhizobacteria. *Agronomic*.9 (8):803-808.

- Rani, B.U. Rajendran, R. K. Suresh (2006). Use of resistant varieties and organic nutrients to manage yellow stem borer in rice. International Rice Research Notes v.31 (2) p.39-41
- Roger, P.A. (1995). Biological N<sub>2</sub> fixation and its management in wetland rice cultivation. Fertilizer Research. 42:261-276.
- Waller, R.A. and D.B. Duncan (1969). Base rule comparison symmetric multiple comparison problem. Amer - stat Assoc. J. 1485-1503.

## استجابة الأرز الهجين مصري واحد إلى مصادر مختلفة من التسميد الأزوتي العضوي والغير عضوي

عبد الفتاح جابر عبد الفتاح ، وليد محمد الخبى ، أحمد محمد الاختيار و بطرس بشرى ميخائيل  
مركز البحوث والتدريب فى الأرز - سخا - كفر الشيخ، معهد بحوث المحاصيل الحقلية - مركز البحوث الزراعية - مصر

أجريت تجربة حقلية بمزرعة مركز البحوث والتدريب فى الأرز بسخا- كفر الشيخ - مصر خلال موسمي صيف ٢٠١٠، ٢٠١١م وذلك لدراسة تأثير مصادر مختلفة للنيتروجين على النمو والمحصول ومكوناته فى مراحل نمو النبات المختلفة حيث تم استخدام الأسمدة غير العضوية مثل اليوريا (٤٦.٥٪ N) وقش الأرز والسماذ البلدي (الحيواني) واللقاح البكتيري Azospirillum SPP (Cerialen) وتم استخدام مخاليط منهم وقد تضمنت التجربة عشرة معاملات كالتالى:-

- (١)- بدون معاملة control (٢)- يوريا (٤٦.٥٪ N) بمقدار ٦٩ كيلوجرام/ فدان
  - (٣)- سماذ بلدي بمقدار ٧طن/ فدان (٤)- قش أرز بمقدار ٢طن / فدان
  - (٥) - لقاح بكتيري Cerialen ٤٠٠ جرام / فدان (٦)- ٤٦ كجم يوريا + ٧طن سماذ بلدي/ فدان
  - (٧)- ٤٦ كجم يوريا + ٢طن قش أرز/ فدان (٨)- ٤٦ كجم يوريا + ٤٠٠ جرام Cerialen
  - (٩)- ٧طن سماذ بلدي + ٤٠٠ جرام Cerialen (١٠)- ٢طن قش أرز + ٤٠٠ جرام Cerialen
- واستخدم فى التجربة تصميم القطاعات كاملة العشوائية فى ثلاثة مكررات ويمكن تلخيص النتائج التي تم الحصول عليها فيما يلى:-

١- أوضحت النتائج أن المعاملات المختلفة للتسميد أدت إلى اختلافات معنوية لمعظم صفات النمو والمحصول و مكوناته موضع الدراسة وكانت أعلى القيم لهذه الصفات فى جميع مراحل النمو المختلفة مع المعاملة ٤٦ كجم يوريا + ٧طن سماذ بلدي (حيواني)/ فدان للصفات - ارتفاع النبات و عدد الأشرطة / الجورة والمادة الجافة المتجمعة محتوى الكلوروفيل بينما كانت أقل قيم للنسبة المئوية للإصابة بالثاقبات عند المعاملة بالسماذ أو قش الأرز مع الخلط باللقاح البكتيري وكانت أقل القيم لهذه الصفات فى القطع غير المعاملة (control)

- وبالنسبة للمحصول ومكوناته :

أوضحت النتائج أن أعلى القيم للصفات المختلفة طول الدالية و عدد الداليات/ الجورة ووزن الدالية (جم) و عدد الحبوب الممتلئة/ الدالية و وزن الألف حبة (جم) و محصول الحبوب (طن/ فدان) { فى حالة المعاملة ٤٦ كجم يوريا + ٧طن سماذ بلدي (حيواني)/ فدان وكذلك أقل القيم لهذه الصفات فى القطع غير المعاملة (control)

\* توصى الدراسة بإمكانية استخدام اليوريا بمقدار ٤٦ كيلوجرام/ فدان مع إضافة ٧طن من السماذ البلدي المتحلل جيدا إلى الفدان لتعظيم إنتاجية وحدة المساحة تحت ظروف محافظة كفر الشيخ

قام بتحكيم البحث

أ.د / احمد ابو النجا قنديل  
كلية الزراعة - جامعة المنصورة  
مركز البحوث الزراعيه

أ.د / محمود رمزى شريف

